

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number
WO 03/093465 A1

- (51) International Patent Classification⁷: C12N 9/64, (74) Common Representative: MAXYGEN ApS; Agern Allé 1, DK-2970 Hørsholm (DK).
C07K 14/745, A61K 38/36, A61P 7/04
- (21) International Application Number: PCT/DK03/00276 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 29 April 2003 (29.04.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/376,679 30 April 2002 (30.04.2002) US
- (71) Applicants (*for all designated States except US*): MAXY-GEN ApS [DK/DK]; Agern Allé 1, DK-2970 Hørsholm (DK). MAXYGEN HOLDINGS LTD. [—/—]; c/o Close Brothers (Cayman) Limited, 103 South Church Street, P.O. Box 1034 GT, Grand Cayman, British West Indies (KY).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): HAANING, Jesper, Mortensen [DK/DK]; Biskop Svanes Vej 2E, DK-3460 Birkerød (DK). ANDERSEN, Kim, Vilbourn [DK/DK]; Karlstads Allé 7, DK-2700 Brønshøj (DK). PEDERSEN, Anders, Hjelholt [DK/DK]; Nybvrovænge 58, DK-2800 Lyngby (DK).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 03/093465 A1

(54) Title: FACTOR VII OR VIIa POLYPEPTIDE VARIANTS

(57) Abstract: The present invention relates to novel polypeptide variants of factor VII (FVII) or factor VIIa (FVIIa) polypeptides, where said variants comprise an amino acid substitution in position 10 and 32 and where said variants further comprise a sugar moiety covalently attached to an introduced *in vivo* N-glycosylation site located outside the Gla domain. Such polypeptide variants are useful in therapy, in particular for the treatment of a variety of coagulation-related disorders, such as trauma.

FACTOR VII OR VIIa POLYPEPTIDE VARIANTS

FIELD OF THE INVENTION

The present invention relates to novel polypeptide variants of factor VII (FVII) or
5 factor VIIa (FVIIa) polypeptides, where said variants comprise an amino acid substitution in
position 10 and 32 and where said variants further comprise a sugar moiety covalently attached
to an introduced *in vivo* N-glycosylation site.

The present invention also relates to use of such polypeptide variants in therapy, in
particular for the treatment of a variety of coagulation-related disorders.

10

BACKGROUND OF THE INVENTION

Blood coagulation is a process consisting of a complex interaction of various blood
components (or factors) that eventually results in a fibrin clot. Generally, the blood components
participating in what has been referred to as the "coagulation cascade" are proenzymes or
15 zymogens, i.e. enzymatically inactive proteins that are converted into an active form by the
action of an activator. One of these coagulation factors is FVII.

FVII is a vitamin K-dependent plasma protein synthesized in the liver and secreted
into the blood as a single-chain glycoprotein with a molecular weight of 53 kDa (Broze &
Majerus, J. Biol. Chem 1980; 255:1242-1247). The FVII zymogen is converted into an
20 activated form (FVIIa) by proteolytic cleavage at a single site, R152-I153, resulting in two
chains linked by a single disulfide bridge. FVIIa in complex with tissue factor (FVIIa complex)
is able to convert both factor IX and factor X into their activated forms, followed by reactions
leading to rapid thrombin production and fibrin formation (Østerud & Rapaport, Proc Natl Acad
Sci USA 1977; 74:5260-5264).

25 FVII undergoes post-translational modifications, including vitamin K-dependent
carboxylation resulting in ten γ -carboxyglutamic acid residues in the N-terminal region of the
molecule. Thus, residue number 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35 shown in SEQ ID NO:2
are γ -carboxyglutamic acids residues in the Gla domain important for FVII activity. Other post-
translational modifications include sugar moiety attachment at two naturally occurring N-
30 glycosylation sites at position 145 and 322, respectively, and at two naturally occurring O-
glycosylation sites at position 52 and 60, respectively.

The gene coding for human FVII (hFVII) has been mapped to chromosome 13 at q34-
qter 9 (de Grouchy et al., Hum Genet 1984; 66:230-233). It contains nine exons and spans 12.8

Kb (O'Hara et al., Proc Natl Acad Sci USA 1987; 84:5158-5162). The gene organisation and protein structure of FVII are similar to those of other vitamin K-dependent procoagulant proteins, with exons 1a and 1b encoding for signal sequence; exon 2 the propeptide and Gla domain; exon 3 a short hydrophobic region; exons 4 and 5 the epidermal growth factor-like domains; and exon 6 through 8 the serine protease catalytic domain (Yoshitake et al., Biochemistry 1985; 24: 3736-3750).

Reports exist on experimental three-dimensional structures of hFVIIa (Pike et al., PNAS. U.S.A., 1999; 96:8925-30 and Kemball-Cook et al., J.Struct.Biol, 1999; 127:213-223); of hFVIIa in complex with soluble tissue factor using X-ray crystallographic methods (Banner et al., Nature, 1996; 380:41 and Zhang et al., J.Mol.Biol, 1999; 285: 2089); and of smaller fragments of hFVII (Muranyi et al., Biochemistry, 1998; 37:10605 and Kao et al., Biochemistry, 1999; 38:7097).

Some protein-engineered variants of FVII have been reported. See, e.g., Dickinson & Ruf, J Bio Chem, 1997;272:19875-19879, Kemball-Cook et al., J Biol Chem, 1998; 273:8516-8521, Bharadwaj et al., J Biol Chem, 1996; 271:30685-30691, Ruf et al., Biochemistry, 1999; 38:1957-1966; WO 99/20767; WO 00/11416; WO 02/22776; WO 02/38162; WO 01/83725; WO 01/58935; US 5,580,560.

Reports exist on expression of FVII in BHK or other mammalian cells (WO 92/15686, WO 91/11514 and WO 88/10295) and co-expression of FVII and kex2 endoprotease in eukaryotic cells (WO 00/28065).

Commercial preparations of human recombinant FVIIa are sold as NovoSeven®. NovoSeven® is indicated for the treatment of bleeding episodes in hemophilia A or B patients. NovoSeven® is the only rhFVIIa for effective and reliable treatment of bleeding episodes available on the market.

An inactive form of FVII in which arginine 152 and/or isoleucine 153 is/are modified has been reported in WO91/1154. These amino acids are located at the activation site. WO 96/12800 describes inactivation of FVIIa by a serine proteinase inhibitor; inactivation by carbamylation of FVIIa at the α -amino acid group I153 has been described by Petersen et al., Eur J Biochem, 1999;261:124-129. The inactivated form is capable of competing with wild-type FVII or FVIIa for binding to tissue factor and inhibiting clotting activity. The inactivated form of FVIIa is suggested to be used for treatment of patients being in hypercoagulable states, such as patients with sepsis, in risk of myocardial infarction or of thrombotic stroke.

A circulating rhFVIIa half-life of 2.3 hours was reported in "Summary Basis for Approval for NovoSeven®", FDA reference number 96-0597. Relatively high doses and frequent administration are necessary to reach and sustain the desired therapeutic or prophylactic effect. As a consequence adequate dose regulation is difficult to obtain and the
5 need of frequent intravenous administrations imposes restrictions on the patient's way of living.

In connection with treatment of uncontrolled bleedings, such as trauma, it is believed that factor VIIa is capable of activating factor X to factor Xa without binding to tissue factor, and this activation reaction is believed to occur primarily on activated blood platelets (Hedner et al. Blood Coagulation & Fibrinolysis, 2000;11;107-111). However, hFVIIa or rhFVIIa has a
10 low activity towards factor X in the absence of tissue factor and, consequently, treatment of uncontrolled bleeding, for example in trauma patients, requires relatively high and multiple doses of hFVIIa or rhFVIIa. Therefore, in order to treat uncontrolled bleedings more efficiently (to minimize blood loss) there is need for improved FVIIa molecules, which possess a high activity toward factor X in the absence of tissue factor. Such improved FVIIa molecules will
15 exhibit a lowered clotting time (or faster action) as compared to rhFVIIa when administered in connection with uncontrolled bleedings.

A molecule with a longer circulation half-life would decrease the number of necessary administrations. Given the association of current the rhFVIIa product with frequent injections, and the potential for obtaining more optimal therapeutic FVIIa levels with concomitant
20 enhanced therapeutic effect, there is a clear need for improved FVII- or FVIIa-like molecules.

One way to increase the circulation half-life of a protein is to ensure that renal clearance of the protein is reduced. This may be achieved by conjugating the protein to a chemical moiety, which is capable of conferring reduced renal clearance to the protein.

Furthermore, attachment of a chemical moiety to the protein or substitution of amino
25 acids exposed to proteolysis may effectively block a proteolytic enzyme from contact leading to proteolytic degradation of the protein. Polyethylene glycol (PEG) is one such chemical moiety that has been used in the preparation of therapeutic protein products. WO 98/32466 suggests that FVII, among many other proteins, may be PEGylated but does not contain any further information in this respect. WO 01/58935 discloses a new strategy for developing FVII or
30 FVIIa molecules having *inter alia* an increased half-life.

As indicated above, another problem in current rhFVIIa treatment is the relative instability of the molecule with respect to proteolytic degradation. Proteolytic degradation is a major obstacle for obtaining a preparation in solution as opposed to a lyophilized product. The advantage of obtaining a stable soluble preparation lies in easier handling for the patient, and, in

the case of emergencies, quicker action, which potentially can become life saving. Attempts to prevent proteolytic degradation by site directed mutagenesis at major proteolytic sites have been disclosed in WO 88/10295.

One object of the present invention is to provide improved FVII or FVIIa molecules
5 (FVII or FVIIa variants) with a longer circulation half-life (thereby decreasing the number of necessary administrations) and which are capable of activating factor X to factor Xa (without binding to tissue factor) more efficiently than hFVIIa or rhFVIIa (thereby being able to treat uncontrolled bleedings, such as a trauma, more efficiently).

Another object of the present invention is to provide improved FVII or FVIIa molecules
10 (FVII or FVIIa variants) with an increased bioavailability (such as an increased Area Under the Curve as compared to rhFVIIa when administered intravenously) and which are capable of activating factor X to factor Xa (without binding to tissue factor) more efficiently than hFVIIa or rhFVIIa (thereby being able to treat uncontrolled bleedings, such as a trauma, more efficiently).

15 These objects are met by the FVII or FVIIa variants provided herein.

BRIEF DISCLOSURE OF THE INVENTION

In its broadest aspect the present invention relates to a FVII or FVIIa polypeptide variant having an amino acid sequence comprising 3-15 amino acid modifications relative to
20 hFVII or hFVIIa having the amino acid sequence shown in SEQ ID NO:2, wherein said amino acid sequence of the variant comprises an amino acid substitution in position 10 and 32 and wherein a sugar moiety is covalently attached to an introduced *in vivo* N-glycosylation site located outside the Gla domain.

Another aspect of the invention relates to a nucleotide sequence encoding the
25 polypeptide variant of the invention.

In a further aspect the invention relates to an expression vector comprising the nucleotide sequence of the invention.

In a still further aspect the invention relates to a host cell comprising the nucleotide sequence of the invention or the expression vector of the invention.

30 In an even further aspect the invention relates to a pharmaceutical composition comprising the polypeptide variant of the invention, and a pharmaceutical acceptable carrier or excipient.

Still another aspect of the invention relates to the polypeptide variant of the invention, or the pharmaceutical composition of the invention, for use as a medicament.

Further aspects of the present invention will be apparent from the below description as well as from the appended claims.

DETAILED DISCLOSURE OF THE INVENTION

5

Definitions

In the context of the present application and invention the following definitions apply:

The term “conjugate” (or interchangeably “conjugated polypeptide”) is intended to indicate a heterogeneous (in the sense of composite or chimeric) molecule formed by the
10 covalent attachment of one or more polypeptide(s) to one or more non-polypeptide moieties such as polymer molecules, lipophilic compounds, sugar moieties or organic derivatizing agents. Preferably, the conjugate is soluble at relevant concentrations and conditions, i.e. soluble in physiological fluids such as blood. Examples of conjugated polypeptides of the invention include glycosylated and/or PEGylated polypeptides.

15 The term “covalent attachment” or “covalently attached” means that the polypeptide variant and the non-polypeptide moiety are either directly covalently joined to one another, or else are indirectly covalently joined to one another through an intervening moiety or moieties, such as a bridge, spacer, or linkage moiety or moieties.

The term “non-polypeptide moiety” is intended to mean a molecule, different from a
20 peptide polymer composed of amino acid monomers and linked together by peptide bonds, which molecule is capable of conjugating to an attachment group of the polypeptide variant of the invention. Preferred examples of such molecules include polymer molecules, sugar moieties, lipophilic compounds or organic derivatizing agents. When used in the context of a conjugated variant of the invention it will be understood that the non-polypeptide moiety is
25 linked to the polypeptide part of the conjugated variant through an attachment group of the polypeptide. As explained above, the non-polypeptide moiety can be directly covalently joined to the attachment group or it can be indirectly covalently joined to the attachment group through an intervening moiety or moieties, such as a bridge spacer or linker moiety or moieties.

A “polymer molecule” is a molecule formed by covalent linkage of two or more
30 monomers, wherein none of the monomers is an amino acid residue, except where the polymer is human albumin or another abundant plasma protein. The term “polymer” may be used interchangeably with the term “polymer molecule”. The term is also intended to cover carbohydrate molecules attached by *in vitro* glycosylation, i.e. a synthetic glycosylation performed *in vitro* normally involving covalently linking a carbohydrate molecule to an

attachment group of the polypeptide variant, optionally using a cross-linking agent. *In vitro* glycosylation is discussed in detail further below.

The term "sugar moiety" is intended to indicate a carbohydrate-containing molecule comprising one or more monosaccharide residues, capable of being attached to the polypeptide
 5 variant (to produce a polypeptide variant conjugate in the form of a glycosylated polypeptide variant) by way of *in vivo* glycosylation. The term "*in vivo* glycosylation" is intended to mean any attachment of a sugar moiety occurring *in vivo*, i.e. during posttranslational processing in a glycosylating cell used for expression of the polypeptide variant, e.g. by way of N-linked and O-linked glycosylation. The exact oligosaccharide structure depends, to a large extent, on the
 10 glycosylating organism in question.

An "N-glycosylation site" has the sequence N-X-S/T/C, wherein X is any amino acid residue except proline, N is asparagine and S/T/C is either serine, threonine or cysteine, preferably serine or threonine, and most preferably threonine. Preferably, the amino acid residue in position +3 relative to the asparagines residue is not a proline residue.

15 An "O-glycosylation site" is the OH-group of a serine or threonine residue.

The term "attachment group" is intended to indicate a functional group of the polypeptide variant, in particular of an amino acid residue thereof or a carbohydrate moiety, capable of attaching a non-polypeptide moiety such as a polymer molecule, a lipophilic molecule, a sugar moiety or an organic derivatizing agent. Useful attachment groups and their
 20 matching non-polypeptide moieties are apparent from the table below.

Attachment group	Amino acid	Examples of non-polypeptide moiety	Conjugation method/-Activated PEG	Reference
-NH ₂	N-terminal, Lys	Polymer, e.g. PEG, with amide or imine group	mPEG-SPA Tresylated mPEG	Shearwater Inc. Delgado et al, critical reviews in Therapeutic Drug Carrier Systems 9(3,4):249-304 (1992)

-COOH	C-terminal, Asp, Glu	Polymer, e.g. PEG, with ester or amide group Carbohydrate moiety	mPEG-Hz <i>In vitro</i> coupling	Shearwater Inc.
-SH	Cys	Polymer, e.g. PEG, with disulfide, maleimide or vinyl sulfone group Carbohydrate moiety	PEG-vinylsul- phone PEG-maleimide <i>In vitro</i> coupling	Shearwater Inc. Delgado et al, critical reviews in Therapeutic Drug Carrier Systems 9(3,4):249-304. (1992)
-OH	Ser, Thr, Lys, OH-	Sugar moiety PEG with ester, ether, carbamate, carbonate	<i>In vivo</i> O-linked glycosylation	
-CONH ₂	Asn as part of an N- glycosyla- tion site	Sugar moiety Polymer, e.g. PEG	<i>In vivo</i> N- glycosylation	
Aromatic residue	Phe, Tyr, Trp	Carbohydrate moiety	<i>In vitro</i> coupling	
-CONH ₂	Gln	Carbohydrate moiety	<i>In vitro</i> coupling	Yan and Wold, Biochemistry, 1984, Jul 31; 23(16): 3759- 65
Aldehyde Ketone	Oxidized oligo- saccharide	Polymer, e.g. PEG, PEG-hydrazide	PEGylation	Andresz et al., 1978, Makromol. Chem. 179:301, WO 92/16555, WO 00/23114

Guanidino	Arg	Carbohydrate moiety	<i>In vitro</i> coupling	Lundblad and Noyes, Chemical Reagents for Protein Modification, CRC Press Inc., Florida, USA
Imidazole ring	His	Carbohydrate moiety	<i>In vitro</i> coupling	As for guanidine

For *in vivo* N-glycosylation, the term "attachment group" is used in an unconventional way to indicate the amino acid residues constituting a N-glycosylation site (with the sequence N-X-S/T/C, wherein X is any amino acid residue except proline, N is asparagine and S/T/C is either serine, threonine or cysteine, preferably serine or threonine, and most preferably threonine). Although the asparagine residue of the N-glycosylation site is the one to which the sugar moiety is attached during glycosylation, such attachment cannot be achieved unless the other amino acid residues of the N-glycosylation site is present.

Accordingly, when the non-polypeptide moiety is a sugar moiety and the conjugation is to be achieved by *in vivo* N-glycosylation, the term "amino acid residue comprising an attachment group for a non-polypeptide moiety" as used in connection with alterations of the amino acid sequence of the polypeptide is to be understood as meaning that one or more amino acid residues constituting an *in vivo* N-glycosylation site are to be altered in such a manner that a functional *in vivo* N-glycosylation site is introduced into the amino acid sequence.

In the present application, amino acid names and atom names (e.g. CA, CB, CD, CG, SG, NZ, N, O, C, etc) are used as defined by the Protein DataBank (PDB) (www.pdb.org) based on the IUPAC nomenclature (IUPAC Nomenclature and Symbolism for Amino Acids and Peptides (residue names, atom names, etc.), *Eur. J. Biochem.*, 138, 9-37 (1984) together with their corrections in *Eur. J. Biochem.*, 152, 1 (1985)).

The term "amino acid residue" is intended to indicate an amino acid residue contained in the group consisting of alanine (Ala or A), cysteine (Cys or C), aspartic acid (Asp or D), glutamic acid (Glu or E), phenylalanine (Phe or F), glycine (Gly or G), histidine (His or H), isoleucine (Ile or I), lysine (Lys or K), leucine (Leu or L), methionine (Met or M), asparagine (Asn or N), proline (Pro or P), glutamine (Gln or Q), arginine (Arg or R), serine (Ser or S), threonine (Thr or T), valine (Val or V), tryptophan (Trp or W), and tyrosine (Tyr or Y) residues.

The terminology used for identifying amino acid positions is illustrated as follows:
I205 indicates that position 205 is occupied by an isoleucine residue in the amino acid sequence shown in SEQ ID NO:2. I205T indicates that the isoleucine residue of position 205 has been substituted with a threonine residue. Alternative substitutions are indicated with a “/”, e.g.
5 I205S/T means an amino acid sequence in which isoleucine in position 205 is substituted with either serine or threonine. Multiple substitutions are indicated with a “+”, e.g. K143N+N145T means a substitution of the lysine residue in position 143 with an asparagine residue and a substitution of the asparagine residue in position 145 with a threonine residue. Insertion of an additional amino acid residue is indicated in the following way: Insertion of a tyrosine residue
10 after A3 (i.e. in position 4) is indicated by A3AY (leading to insertion of a tyrosine residue in position 4). A deletion of an amino acid residue is indicated by an asterix. For example, deletion of a valine residue in position 172 is indicated by V172*. Simultaneous insertion and substitution are indicated in the following way: Substitution of an alanine residue in position 175 with a threonine residue followed by insertion of a leucine residue after position 175 is
15 indicated A175TL.

Unless otherwise indicated, the numbering of amino acid residues made herein is made relative to the amino acid sequence of the hFVII/hFVIIa polypeptide (SEQ ID NO:2).

The term “differs from” as used in connection with specific mutations is intended to allow for additional differences being present apart from the specified amino acid difference.
20 For instance, in addition to the introduction of *in vivo* N-glycosylation sites (located outside the Gla domain), the polypeptide may comprise other modifications that are not related to introduction of such amino acid residues. In a similar way, in addition to the modifications performed in the Gla domain aiming at increasing the phospholipid membrane binding affinity, the polypeptide may contain other modifications that are not necessarily related to this effect.

25 Thus, in addition to the amino acid modifications disclosed herein, it will be understood that the amino acid sequence of the polypeptide variant of the invention may, if desired, contain other alterations, i.e. other substitutions, insertions or deletions. These may, for example, include truncation of the N- and/or C-terminus by one or more amino acid residues (e.g. by 1-10 amino acid residues), or addition of one or more extra residues at the N- and/or C-
30 terminus, e.g. addition of a methionine residue at the N-terminus or introduction of a cysteine residue near or at the C-terminus, as well as “conservative amino acid substitutions”, i.e. substitutions performed within groups of amino acids with similar characteristics, e.g. small amino acids, acidic amino acids, polar amino acids, basic amino acids, hydrophobic amino acids and aromatic amino acids.

Examples of such conservative substitutions are shown in the below table.

1	Alanine (A)	Glycine (G)	Serine (S)	Threonine (T)
2	Aspartic acid (D)	Glutamic acid (E)		
3	Asparagine (N)	Glutamine (Q)		
4	Arginine (R)	Histidine (H)	Lysine (K)	
5	Isoleucine (I)	Leucine (L)	Methionine (M)	Valine (V)
6	Phenylalanine (F)	Tyrosine (Y)	Tryptophan (W)	

Still other examples of additional modifications are disclosed in the section entitled
 5 *"Other modifications outside the Gla domain"* below.

The term "nucleotide sequence" is intended to indicate a consecutive stretch of two or more nucleotide molecules. The nucleotide sequence may be of genomic, cDNA, RNA, semisynthetic, synthetic origin, or any combinations thereof.

The term "polymerase chain reaction" or "PCR" generally refers to a method for
 10 amplification of a desired nucleotide sequence *in vitro*, as described, for example, in US 4,683,195. In general, the PCR method involves repeated cycles of primer extension synthesis, using oligonucleotide primers capable of hybridising preferentially to a template nucleic acid.

The term "vector" refers to a plasmid or other nucleotide sequences that are capable of replicating within a host cell or being integrated into the host cell genome, and as such, are
 15 useful for performing different functions in conjunction with compatible host cells (a vector-host system): to facilitate the cloning of the nucleotide sequence, i.e. to produce usable quantities of the sequence, to direct the expression of the gene product encoded by the sequence and to integrate the nucleotide sequence into the genome of the host cell. The vector will contain different components depending upon the function it is to perform.

20 "Cell", "host cell", "cell line" and "cell culture" are used interchangeably herein and all such terms should be understood to include progeny resulting from growth or culturing of a cell.

"Transformation" and "transfection" are used interchangeably to refer to the process of introducing DNA into a cell.

25 "Operably linked" refers to the covalent joining of two or more nucleotide sequences, by means of enzymatic ligation or otherwise, in a configuration relative to one another such that the normal function of the sequences can be performed. For example, the nucleotide sequence encoding a presequence or secretory leader is operably linked to a nucleotide sequence coding

for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide: a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the
5 nucleotide sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, then synthetic oligonucleotide adaptors or linkers are used, in conjunction with standard recombinant DNA methods.

In the context of the present invention the terms "modification" or "amino acid
10 modification" is intended to cover replacement of an amino acid side chain, substitution of an amino acid residue, deletion of an amino acid residue and/or insertion of an amino acid residue.

The terms "mutation" and "substitution" are used interchangeably herein.

The term "introduce" refers to introduction of an amino acid residue by substitution of an existing amino acid residue or, alternatively, by insertion of an additional amino acid
15 residue.

The term "remove" refers to removal of an amino acid residue by substitution of the amino acid residue to be removed by another amino acid residue or, alternatively, by deletion (without substitution) of the amino acid residue to be removed

The term "FVII" or "FVII polypeptide" refers to a FVII molecule provided in single
20 chain form. One example of a FVII polypeptide is wild-type human FVII (hFVII) shown in SEQ ID NO:2. It should be understood, however, that the term "FVII polypeptide" also covers hFVII-like molecules, such as fragments or variants of SEQ ID NO:2, in particular variants where the sequence comprises at least one, such as 1-15, e.g., 1-10, amino acid modifications as compared to SEQ ID NO:2.

25 The term "FVIIa" or "FVIIa polypeptide" refers to a FVIIa molecule provided in its activated two-chain form. When the amino acid sequence of SEQ ID NO:2 is used to describe the amino acid sequence of FVIIa it will be understood that the peptide bond between R152 and I153 of the single-chain form has been cleaved, and that one of the chains comprises amino acid residues 1-152, the other chain amino acid residues 153-406.

30 The terms "rFVII" and "rFVIIa" refer to FVII and FVIIa polypeptides produced by recombinant techniques.

The terms "hFVII" and "hFVIIa" refer to human wild-type FVII and FVIIa, respectively, having the amino acid sequence shown in SEQ ID NO:2

The terms "rhFVII" and "rhFVIIa" refer to human wild-type FVII and FVIIa, having the amino acid sequence shown in SEQ ID NO:2, produced by recombinant means. An example of rhFVIIa is NovoSeven®.

When used herein, the term "Gla domain" is intended to cover amino acid residues no. 1 to 45 of SEQ ID NO:2.

Accordingly, the term "located outside the Gla domain" covers amino acid residue no. 46-406 of SEQ ID NO:2.

The abbreviations "TF" and "TFPI" mean Tissue Factor and Tissue Factor Pathway Inhibitor, respectively.

The term "protease domain" is used about residues 153-406 counted from the N-terminus.

The term "catalytic site" is used to mean the catalytic triad consisting of S344, D242 and H193 of the polypeptide variant.

The term "parent" is intended to indicate the molecule to be modified/improved in accordance with the present invention. Although the parent polypeptide to be modified by the present invention may be any FVII or FVIIa polypeptide, and thus be derived from any origin, e.g. a non-human mammalian origin, it is preferred that the parent polypeptide is hFVII or hFVIIa.

A "variant" is a polypeptide, which differs in one or more amino acid residues from its parent polypeptide, normally in 3-15 amino acid residues (e.g. in 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid residues), such as in 3-10 amino acid residues, e.g. in 3-8 or 3-5 amino acid residues. In other words, a "variant" typically contains 3-15 amino acid modifications (for example 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid modifications), such as 3-10 amino acid modifications, e.g. 3-8 or 3-5 amino acid modifications relative to the parent polypeptide. In the present context, the term "modification" encompasses insertions, deletions, substitutions and combinations thereof. It will be understood that a polypeptide variant according to the present invention will be modified in at least three positions, namely in at least position 10 and 32 (located in the Gla domain) and in at least one position located outside the Gla domain, where said at least one modification creates an *in vivo* N-glycosylation site.

The term "clotting activity" is used to mean the activity measured in the "Clotting Assay" described herein. In order to exhibit "clotting activity" a variant of the invention, in its activated form, should have at least 10% of the clotting activity of rhFVIIa when assayed in the "Clotting Assay" described herein. In a preferred embodiment of the invention the variant, in its activated form, has at least 20% of the clotting activity of rhFVIIa, such as at least 30%, e.g. at

least 40%, more preferably at least 50%, such as at least 60%, e.g. at least 70%, even more preferably at least 80%, such as at least 90% of the clotting activity of rhFVIIa when assayed in the "Clotting Assay" described herein. In an interesting embodiment the variant, in its activated form, has substantially the same clotting activity as rhFVIIa, such as a clotting activity of 75-
5 125% of the clotting activity of rhFVIIa.

The term "amidolytic activity" is used to mean the activity measured in the "Amidolytic Assay" described herein. In order to exhibit "amidolytic activity" a variant of the invention, in its activated form, should have at least 10% of the amidolytic activity of rhFVIIa when assayed in the "Amidolytic Assay" described herein. In a preferred embodiment of the
10 invention the variant, in its activated form, has at least 20% of the amidolytic activity of rhFVIIa, such as at least 30%, e.g. at least 40%, more preferably at least 50%, such as at least 60%, e.g. at least 70%, even more preferably at least 80%, such as at least 90% of the amidolytic activity of rhFVIIa when assayed in the "Amidolytic Assay" described herein. In an interesting embodiment the variant, in its activated form, has substantially the same amidolytic
15 activity as rhFVIIa, such as an amidolytic activity of 75-125% of the amidolytic activity of rhFVIIa.

In the present context, the term "activity" is also used in connection with the variants capability of activating FX to FXa. This activity is also denoted "FX activation activity" or "FXa generation activity".

20 The term "increased FX activation activity" or "increased FXa generation activity" is used to indicate that a variant of the invention, in its activated form, has a statistically significantly increased capability to activate FX to FXa as compared to rhFVIIa or. To what extent a variant of the invention (in its activated form) has an increased FX activation activity may conveniently be determined in the "TF-independent Factor X Activation Assay" described
25 herein.

The term "stronger clot" or "increased clot strength" is used to indicate that the strength of the clot generated by the polypeptide variant is statistically significantly increased relative to that generated by rhFVIIa as determined under comparable conditions. This effect may be determined as the Area Under the Curve (AUC_{throm}) generated by the variant of the
30 invention, in its activated form, when assayed in the "Thrombogram Assay" disclosed herein. In a similar way, the term "increased AUC_{throm} " is used to indicate that the Area Under the Curve generated by the variant (in its activated form) is statistically significantly increased relative to that generated by rhFVIIa as determined under comparable conditions and when measured in the "Thrombogram Assay" described herein

The term " T_{\max} " is used about the time it takes to obtain the maximum thrombin activity level in the "Thrombogram Assay".

The term "immunogenicity" as used in connection with a given substance is intended to indicate the ability of the substance to induce a response from the immune system. The immune response may be a cell or antibody mediated response (see, e.g., Roitt: Essential Immunology (8th Edition, Blackwell) for further definition of immunogenicity). Normally, reduced antibody reactivity will be an indication of reduced immunogenicity. The reduced immunogenicity may be determined by use of any suitable method known in the art, e.g. *in vivo* or *in vitro*.

The term "functional *in vivo* half-life" is used in its normal meaning, i.e. the time at which 50% of the biological activity of the polypeptide is still present in the body/target organ, or the time at which the activity of the polypeptide is 50% of the initial value.

As an alternative to determining functional *in vivo* half-life, "serum half-life" may be determined, i.e. the time at which 50% of the polypeptide circulates in the plasma or bloodstream prior to being cleared. Determination of serum half-life is often more simple than determining the functional *in vivo* half-life and the magnitude of serum half-life is usually a good indication of the magnitude of functional *in vivo* half-life. Alternatively terms to serum half-life include "plasma half-life", "circulating half-life", "serum clearance", "plasma clearance" and "clearance half-life". The polypeptide is cleared by the action of one or more of the reticuloendothelial systems (RES), kidney, spleen or liver, by tissue factor, SEC receptor or other receptor mediated elimination, or by specific or unspecific proteolysis. Normally, clearance depends on size (relative to the cutoff for glomerular filtration), charge, attached carbohydrate chains, and the presence of cellular receptors for the protein. The functionality to be retained is normally selected from procoagulant, proteolytic or receptor binding activity. The functional *in vivo* half-life and the serum half-life may be determined by any suitable method known in the art.

The term "increased" as used about the functional *in vivo* half-life or serum half-life is used to indicate that the relevant half-life of the polypeptide variant is statistically significantly increased relative to that of rhFVIIa as determined under comparable conditions (typically determined in an experimental animal, such as rats, rabbits, pigs or monkeys).

The term " AUC_{iv} " or "Area Under the Curve when administered intravenously" is used in its normal meaning, i.e. as the area under the activity in serum-time curve, where the polypeptide variant has been administered intravenously, in particular when administered intravenously in rats. Typically, the activity measured is the "clotting activity" as defined

hereinbefore. Once the experimental activity-time points have been determined, the AUC_{iv} may conveniently be calculated by a computer program, such as GraphPad Prism 3.01.

It will be understood that in order to make a direct comparison between the AUC_{iv} -values of different molecules (e.g. between the variants of the invention and a reference molecule such as rhFVIIa) the same amount of activity should be administered. Consequently, the AUC_{iv} -values are typically normalized (i.e. corrected for differences in the injected dose) and expressed as $AUC_{iv}/\text{dose administered}$.

The term "reduced sensitivity to proteolytic degradation" is primarily intended to mean that the polypeptide variant has reduced sensitivity to proteolytic degradation in comparison to hFVIIa or rhFVIIa as determined under comparable conditions. Preferably, the proteolytic degradation is reduced by at least 10% (e.g. by 10-25% or by 10-50%), such as at least 25% (e.g. by 25-50%, by 25-75% or by 25-100%), more preferably by at least 35%, such as at least 50%, (e.g. by 50-75% or by 50-100%) even more preferably by at least 60%, such as by at least 75% (e.g. by 75-100%) or even at least 90%. Most preferably, the proteolytic degradation is reduced by at least 100%.

The term "renal clearance" is used in its normal meaning to indicate any clearance taking place by the kidneys, e.g. by glomerular filtration, tubular excretion or degradation in the tubular cells. Renal clearance depends on physical characteristics of the polypeptide, including size (diameter), hydrodynamic volume, symmetry, shape/rigidity, and charge. Normally, a molecular weight of about 67 kDa is considered to be a cut-off-value for renal clearance. Renal clearance may be established by any suitable assay, e.g. an established *in vivo* assay. Typically, renal clearance is determined by administering a labelled (e.g. radiolabelled or fluorescence labelled) polypeptide to a patient and measuring the label activity in urine collected from the patient. Reduced renal clearance is determined relative to a corresponding reference polypeptide, e.g. rhFVIIa, under comparable conditions. Preferably, the renal clearance rate of the polypeptide variant is reduced by at least 50%, preferably by at least 75%, and most preferably by at least 90% compared to rhFVIIa.

The terms "at least 25% of its side chain exposed to the surface of the molecule" and "at least 50% of its side chain exposed to the surface of the molecule" are defined with reference to Example 1, where the calculations, etc. are described in detail.

It should be noted that when the terms "at least 25% of its side chain exposed to the surface of the molecule" and "at least 50% of its side chain exposed to the surface of the molecule" are used in connection with introduction of an *in vivo* N-glycosylation site these terms refer to the surface accessibility of the amino acid side chain in the position where the

sugar moiety is actually attached. In many cases it will be necessary to introduce a serine or a threonine residue in position +2 relative to the asparagine residue to which the sugar moiety is actually attached (unless, of course, this position is already occupied by a serine or a threonine residue) and these positions, where the serine or threonine residues are introduced, are allowed to be buried, i.e. to have less than 25% or 50% of their side chains exposed to the surface of the molecule.

The terms "tissue factor binding site", "active site region" and "ridge of the active site binding cleft" are defined with reference to Example 1 herein, wherein the above-mentioned sites/regions are determined.

10 The term "hydrophobic amino acid residue" includes the following amino acid residues: isoleucine (I), leucine (L), methionine (M), valine (V), phenylalanine (F), tyrosine (Y) and tryptophan (W).

The term "negatively charged amino acid residue" includes the following amino acid residues: Aspartic acid (D) and glutamic acid (E).

15 The term "positively charged amino acid residue" includes the following amino acid residues: Lysine (K), arginine (R) and histidine (H).

Variants of the invention

In its broadest aspect the present invention relates to a FVII or FVIIa polypeptide variant having an amino acid sequence comprising 3-15 amino acid modifications relative to hFVII or hFVIIa having the amino acid sequence shown in SEQ ID NO:2, wherein said amino acid sequence of the variant comprises an amino acid substitution in position 10 and 32 and wherein a sugar moiety is covalently attached to an introduced *in vivo* N-glycosylation site located outside the Gla domain.

25 The modifications performed in the two above-indicated regions of the parent FVII polypeptide serve the following purposes:

The modifications performed in positions located outside the Gla domain (introduction of *in vivo* N-glycosylation site(s)) are preferably of such nature that the AUC_{iv}, the functional *in vivo* half-life and/or the serum half-life of the resulting variant is increased as compared to rhFVIIa.

30 The modifications performed in the Gla domain of the parent polypeptide are preferably of such nature that an increased phospholipid membrane binding affinity of the resulting molecule is achieved and/or of such nature the resulting molecule has an improved capability to activate FX to FXa and/or of such nature that a stronger clot is formed.

Without being limited by any particular theory, it is presently believed that the enhanced membrane affinity results in a higher local concentration of the activated polypeptide variants in close proximity to the other coagulation factors, particularly FX. Thus, the rate of activation of FX to FXa will be higher, simply due to a higher molar ratio of the activated FVII variant to FX. The increased activation rate of FX then results in a higher amount of active thrombin, and thus a higher rate of cross-linking of fibrin.

Thus, in preferred embodiments of the invention the parent FVII or FVIIa polypeptide has been modified so that the resulting activated polypeptide variant has (as compared to rhFVIIa):

- i) an increased bioavailability (AUC_{iv}) and an increased phospholipid membrane binding affinity;
- ii) an increased bioavailability (AUC_{iv}) and an increased capability to activate FX to FXa;
- iii) an increased bioavailability (AUC_{iv}) and is capable of generating a stronger clot (increased AUC_{throm});
- iv) an increased bioavailability (AUC_{iv}) and a reduced T_{max} ;
- v) an increased functional *in vivo* half-life and an increased phospholipid membrane binding affinity;
- vi) an increased functional *in vivo* half-life and an increased capability to activate FX to FXa;
- vii) an increased functional *in vivo* half-life and is capable of generating a stronger clot (increased AUC_{throm});
- viii) an increased functional *in vivo* half-life and a reduced T_{max} ;
- ix) an increased serum half-life and an increased phospholipid membrane binding affinity;
- x) an increased serum half-life and an increased capability to activate FX to FXa;
- xi) an increased serum half-life and is capable of generating a stronger clot (increased AUC_{throm}); and/or
- xii) an increased serum half-life and a reduced T_{max} .

Consequently, medical treatment with a polypeptide variant according to the invention offers a number of advantages over the currently available rhFVIIa compound (NovoSeven®), such as administration of lower dosage, longer duration between injections, increased clot strength and/or faster action.

Thus, preferred variants of the invention are such variants which, in their activated forms and when compared to rhFVIIa, generates in increased Area Under the Curve when administered intravenously (AUC_{iv}), in particular when administered intravenously in rats. More particularly, variants of the present invention, which are preferred, are such variants
5 where the ratio between the AUC_{iv} of said variant, in its activated form, and the AUC_{iv} of rhFVIIa is at least 1.25, such as at least 1.5, e.g. at least 1.75, more preferably at least 2, such as at least 3, even more preferably at least 4, such as at least 5, in particular when administered (intravenously) in rats.

This effect may in turn (but do not necessarily do so) correspond to an increased
10 functional *in vivo* half-life and/or an increased serum half-life as compared to rhFVIIa. Accordingly, in another preferred embodiment of the invention, the ratio between the functional *in vivo* half-life or the serum half-life for the variant, in its activated form, and the functional *in vivo* half-life or the serum half-life for rhFVIIa is at least 1.25. More preferably, the ratio between the relevant half-life for the variant, in its activated form, and the relevant half-life for
15 hFVIIa or rhFVIIa is at least 1.5, such as at least 1.75, e.g. at least 2, even more preferably at least 3, such as at least 4, e.g. at least 5.

As will be understood, the variants of the invention also possess, in addition to the above-mentioned functionality (i.e. increased AUC_{iv} , increased functional *in vivo* half-life and/or increased serum half-life), an increased phospholipid membrane binding affinity as
20 compared to rhFVII, an an increased capability to activate FX to Fxa, a capability of generating a stronger clot (increased AUC_{throm}) and/or a reduced T_{max} .

Thus, in one preferred embodiment of the invention, the polypeptide variant has (in addition to an increased AUC_{iv} , increased functional *in vivo* half-life and/or increased serum half-life) an increased phospholipid membrane binding affinity relative to rhFVIIa. Membrane
25 binding affinity may be measured by methods known in the art, such as by the Biacore® assay described in K. Nagata and H. Handa (Eds.), Real-Time Analysis of Biomolecular Interactions, Springer-Verlag, Tokyo, 2000, Chapter 6 entitled "Lipid-Protein Interactions". Alternatively, the membrane binding affinity may be measured as described in Example 1 in WO 99/20767.

In another preferred embodiment of the invention, the polypeptide variant (in addition
30 to an increased AUC_{iv} , increased functional *in vivo* half-life and/or increased serum half-life), has an increased FX activation activity as compared to rhFVIIa, in particular when assayed in a TF-independent assay, such as the "TF-independent Factor X Activation Assay" disclosed herein. More particularly, it is preferred that the ratio between the FX activation activity of the polypeptide variant, in its activated form, and the FX activation activity of rhFVIIa is at least

1.25 when assayed in the "TF-independent Factor X Activation Assay" disclosed herein. More preferably, the ratio between the FX activation activity of the variant, in its activated form, and the FX activation activity of rhFVIIa is at least 1.5, such as at least 1.75, e.g. at least 2, even more preferably at least 3, such as at least 4, e.g. at least 5, still more preferably at least 6, such as at least 7, e.g. at least 8, most preferably at least 9, such as at least 10, when assayed in the "TF-independent Factor X Activation Assay" described herein.

In still another preferred embodiment of the invention, the polypeptide variant is (in addition to an increased AUC_{iv} , increased functional *in vivo* half-life and/or increased serum half-life), in its activated form, capable of generating a stronger clot as compared to rhFVIIa. This effect may be determined in the "Thrombogram Assay" described herein as an increase in the Area Under the Curve (AUC_{throm}). The AUC_{throm} is also sometimes denoted "total thrombin work" and constitutes a measure for the strength of the clot formed. More particularly, it is preferred that the ratio between the AUC_{throm} , generated by the variant in its activated form, and the AUC_{throm} generated by rhFVIIa is at least 1.15 when assayed in the "Thrombogram Assay" described herein. More preferably, the ratio is at least 1.2, such as at least 1.25, e.g. at least 1.3, even more preferably at least 1.4, such as at least 1.5, e.g. at least 1.6, most preferably at least 1.7, such as at least 1.8, e.g. at least 1.9 or at least 2.

In even another preferred embodiment of the invention the polypeptide variant has (in addition to an increased AUC_{iv} , increased functional *in vivo* half-life and/or increased serum half-life), in its activated form, a faster action. This effect may be determined in the "Thrombogram Assay" described herein as a reduction in the time needed to reach maximum thrombin level (T_{max}). Accordingly, preferred variants are such variants where the ratio between T_{max} for the variant, in its activated form, and T_{max} for rhFVIIa is at the most 0.95 when assayed in the "Thrombogram Assay" described herein. Preferably, the ratio is at the most 0.9, such as at the most 0.8, e.g. at the most 0.7, more preferably at the most 0.6, such as at the most 0.5.

Introduction of *in vivo* N-glycosylation sites located outside the Gla domain

A number of suitable modifications leading to an increase in AUC_{iv} , functional *in vivo* half-life and/or serum half-life is disclosed in WO 01/58935. The variants disclosed in WO 01/58935 are the result of a generally new strategy for developing improved FVII or FVIIa molecules, which may also be used for the parent FVII or FVIIa polypeptide of the present invention.

The position to be modified is preferably selected from a part of the FVII or FVIIa molecule that is located outside the tissue factor binding site, and/or outside the active site

region, and/or outside the ridge of the active site binding cleft. These sites/regions are identified in Example 1 herein. It should be emphasized, however, that in certain situations, e.g. in case an inactivated polypeptide variant is desired, it may be advantageous to perform modifications in or close to such regions. For example, it is contemplated that one or more *in vivo* N-glycosylation sites may advantageously be introduced in the active site region or at the ridge of the active site binding cleft of the FVII or FVIIa molecule. The active site region, the tissue factor binding site and the ridge of the active site binding cleft are defined in Example 1 herein and are constituted by the following residues:

I153, Q167, V168, L169, L170, L171, Q176, L177, C178, G179, G180, T181, V188, V189, S190, A191, A192, H193, C194, F195, D196, K197, I198, W201, V228, I229, I230, P231, S232, T233, Y234, V235, P236, G237, T238, T239, N240, H241, D242, I243, A244, L245, L246, V281, S282, G283, W284, G285, Q286, T293, T324, E325, Y326, M327, F328, D338, S339, C340, K341, G342, D343, S344, G345, G346, P347, H348, L358, T359, G360, I361, V362, S363, W364, G365, C368, V376, Y377, T378, R379, V380, Q382, Y383, W386, L387, L400 and F405 (active site region);

L13, K18, F31, E35, R36, L39, F40, I42, S43, S60, K62, D63, Q64, L65, I69, C70, F71, C72, L73, P74, F76, E77, G78, R79, E82, K85, Q88, I90, V92, N93, E94, R271, A274, F275, V276, R277, F278, R304, L305, M306, T307, Q308, D309, Q312, Q313, E325 and R379 (tissue factor binding site); and

N173, A175, K199, N200, N203, D289, R290, G291, A292, P321 and T370 (the ridge of the active site binding cleft).

The total number of amino acid residues to be modified outside the Gla domain in the parent FVII or FVIIa polypeptide (as compared to the amino acid sequence shown in SEQ ID NO:2) will typically not exceed 10. Preferably, the FVII or FVIIa variant comprises an amino acid sequence which differs in 1-10 amino acid residues from amino acid residues 46-406 shown in SEQ ID NO:2, typically in 1-8 or in 2-8 amino acid residues, e.g. in 1-5 or in 2-5 amino acid residues, such as in 1-4 or in 1-3 amino acid residues, e.g. in 1, 2 or 3 amino acid residues from amino acid residues 46-406 shown in SEQ ID NO:2.

Thus, the polypeptide variant of the invention may contain 1-10 (additional or introduced) *in vivo* N-glycosylation sites, typically 1-8 or 2-8 (additional or introduced) *in vivo* N-glycosylation sites, preferably 1-5 or 2-5 (additional or introduced) *in vivo* N-glycosylation sites, such as 1-4 or 1-3 (additional or introduced) *in vivo* N-glycosylation sites, e.g. 1, 2 or 3 (additional or introduced) *in vivo* N-glycosylation sites. Analogously, the polypeptide variant of the invention may contain 1-10 (additional or introduced) sugar moieties, typically 1-8 or 2-8

(additional or introduced) sugar moieties, preferably 1-5 or 2-5 (additional or introduced) sugar moieties, such as 1-4 or 1-3 (additional or introduced) sugar moieties, e.g. 1, 2 or 3 (additional or introduced) sugar moieties. It will be understood that the introduced sugar moiety/moieties will be covalently attached to the introduced *in vivo* N-glycosylation site(s)

5 When used in the present context, the term "naturally occurring glycosylation site" covers the glycosylation sites at positions N145, N322, S52 and S60. In a similar way, the term "naturally occurring *in vivo* O-glycosylation site" includes the positions S52 and S60, whereas the term "naturally occurring *in vivo* N-glycosylation site" includes positions N145 and N322.

It will be understood that in order to prepare a polypeptide variant, wherein the
10 polypeptide variant comprises one or more sugar moieties covalently attached to one or more *in vivo* N-glycosylation sites, the polypeptide variant must be expressed in a host cell capable of attaching sugar (oligosaccharide) moieties at the glycosylation site(s) or alternatively subjected to *in vitro* glycosylation. Examples of glycosylating host cells are given in the section further below entitled "*Coupling to a sugar moiety*".

15 Examples of positions, wherein the *in vivo* N-glycosylation sites may be introduced include, but is not limited to, positions comprising an amino acid residue having an amino acid residue having at least 25% of its side chain exposed to the surface (as defined in Example 1 herein), such as in a position comprising an amino acid residue having at least 50% of its side chain exposed to the surface (as defined in Example 1 herein). In general, it is preferred that the
20 *in vivo* N-glycosylation site is introduced by substitution, although insertion is also contemplated. The position is preferably selected from a part of the molecule that is located outside the tissue factor binding site and/or the active site region and/or outside the ridge of the active site cleft. These sites/regions are identified in Example 1 herein. It should be understood that when the term "at least 25% (or at least 50%) of its side chain exposed to the surface" is used in
25 connection with introduction of an *in vivo* N-glycosylation site this term refers to the surface accessibility of the amino acid side chain in the position where the sugar moiety is actually attached. In many cases it will be necessary to introduce a serine or a threonine residue in position +2 relative to the asparagine residue to which the sugar moiety is actually attached (unless, of course, this position is already occupied by a serine or a threonine residue) and these
30 positions, where the serine or threonine residues are introduced, are allowed to be buried, i.e. to have less than 25% of their side chains exposed to the surface.

Specific and preferred examples of such substitutions creating an *in vivo* N-glycosylation site include a substitution selected from the group consisting of A51N, G58N, T106N, K109N, G124N, K143N+N145T, A175T, I205S, I205T, V253N, T267N,

T267N+S269T, S314N+K316S, S314N+K316T, R315N+V317S, R315N+V317T, K316N+G318S, K316N+G318T, G318N, D334N and combinations thereof. More preferably, the *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of A51N, G58N, T106N, K109N, G124N, K143N+N145T, A175T, I205T, V253N,
 5 T267N+S269T, S314N+K316T, R315N+V317T, K316N+G318T, G318N, D334N and combinations thereof. Even more preferably, the *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of T106N, A175T, I205T, V253N, T267N+S269T and combinations thereof, in particular I205T.

In one embodiment, only one *in vivo* N-glycosylation site has been introduced by
 10 substitution. In another embodiment, two or more (such as two) *in vivo* N-glycosylation sites have been introduced by substitution. Examples of preferred substitutions creating two *in vivo* N-glycosylation sites include substitutions selected from the group consisting of A51N+G58N, A51N+T106N, A51N+K109N, A51N+G124N, A51N+K143N+N145T, A51N+A175T, A51N+I205T, A51N+V253N, A51N+T267N+S269T, A51N+S314N+K316T,
 15 A51N+R315N+V317T, A51N+K316N+G318T, A51N+G318N, A51N+D334N, G58N+T106N, G58N+K109N, G58N+G124N, G58N+K143N+N145T, G58N+A175T, G58N+I205T, G58N+V253N, G58N+T267N+S269T, G58N+S314N+K316T, G58N+R315N+V317T, G58N+K316N+G318T, G58N+G318N, G58N+D334N, T106N+K109N, T106N+G124N, T106N+K143N+N145T, T106N+A175T, T106N+I205T,
 20 T106N+V253N, T106N+T267N+S269T, T106N+S314N+K316T, T106N+R315N+V317T, T106N+K316N+G318T, T106N+G318N, T106N+D334N, K109N+G124N, K109N+K143N+N145T, K109N+A175T, K109N+I205T, K109N+V253N, K109N+T267N+S269T, K109N+S314N+K316T, K109N+R315N+V317T, K109N+K316N+G318T, K109N+G318N, K109N+D334N, G124N+K143N+N145T,
 25 G124N+A175T, G124N+I205T, G124N+V253N, G124N+T267N+S269T, G124N+S314N+K316T, G124N+R315N+V317T, G124N+K316N+G318T, G124N+G318N, G124N+D334N, K143N+N145T+A175T, K143N+N145T+I205T, K143N+N145T+V253N, K143N+N145T+T267N+S269T, K143N+N145T+S314N+K316T, K143N+N145T+R315N+V317T, K143N+N145T+K316N+G318T, K143N+N145T+G318N,
 30 K143N+N145T+D334N, A175T+I205T, A175T+V253N, A175T+T267N+S269T, A175T+S314N+K316T, A175T+R315N+V317T, A175T+K316N+G318T, A175T+G318N, A175T+D334N, I205T+V253N, I205T+T267N+S269T, I205T+S314N+K316T, I205T+R315N+V317T, I205T+K316N+G318T, I205T+G318N, I205T+D334N,

V253N+T267N+S269T, V253N+S314N+K316T, V253N+R315N+V317T, V253N+K316N+G318T, V253N+G318N, V253N+D334N, T267N+S269T+S314N+K316T, T267N+S269T+R315N+V317T, T267N+S269T+K316N+G318T, T267N+S269T+G318N, T267N+S269T+D334N, S314N+K316T+R315N+V317T, S314N+K316T+G318N, 5 S314N+K316T+D334N, R315N+V317T+K316N+G318T, R315N+V317T+G318N, R315N+V317T+D334N and G318N+D334N. More preferably, the substitutions are selected from the group consisting of T106N+A175T, T106N+I205T, T106N+V253N, T106N+T267N+S269T, A175T+I205T, A175T+V253N, A175T+T267N+S269T, I205T+V253N, I205T+T267N+S269T and V253N+T267N+S269T, even more preferably from 10 the group consisting of T106N+I205T, T106N+V253N and I205T+T267N+S269T.

In an even further embodiment, three or more (such as three) *in vivo* N-glycosylation sites have been introduced by substitution. Examples of preferred substitutions creating three *in vivo* N-glycosylation sites include substitutions selected from the group consisting of I205T+V253N+T267N+S269T and T106N+I205T+V253N.

15 As discussed above, it is preferred that the *in vivo* N-glycosylation site is introduced in a position which does neither form part of the tissue factor binding site nor form part of the active site region and the ridge of the active site binding cleft as defined herein. It is envisaged that such glycosylation variants will primarily belong to the class of active polypeptide variants as defined hereinbefore.

20 It will be understood that as an alternative to introduction of *in vivo* N-glycosylation sites in the above-discussed positions one may introduce (either by substitution or by insertion) cysteine residues in the same positions, where the introduced cysteine residue is then subsequently covalently attached to a non-polypeptide moiety, such as PEG, in particular mPEG. Thus, examples of positions, wherein a cysteine residue may be introduced include, but 25 is not limited to, positions comprising an amino acid residue having an amino acid residue having at least 25% of its side chain exposed to the surface (as defined in Example 1 herein), such as in a position comprising an amino acid residue having at least 50% of its side chain exposed to the surface (as defined in Example 1 herein). The position is preferably selected from a part of the molecule that is located outside the tissue factor binding site and/or the active 30 site region and/or outside the ridge of the active site cleft. These sites/regions are identified in Example 1 herein. Thus, the above disclosure concerning introduction of *in vivo* N-glycosylation sites applies mutatis mutandis to introduction of cysteine residues.

It will be understood that the modifications in positions located outside the Gla domain discussed in the above section should be combined with one or more modifications in the Gla domain (see the section entitled "*Modifications in the Gla domain*" below).

5 *Modifications in the Gla domain*

As will be understood the variants of the present invention comprises, in addition to at least one introduced *in vivo* N-glycosylation site located outside the Gla domain (cf. above), at least two substitutions in the Gla domain, namely a substitution in position 10 and position 32.

A number of suitable modifications leading to increased phospholipid membrane
10 binding affinity is disclosed in WO 99/20767 and WO 00/66753.

In a preferred embodiment of the invention the substitution in position 10 is P10Q. In another preferred embodiment of the invention the substitution in position 32 is K32E. In a particularly preferred embodiment of the invention the variant comprises the following substitutions P10Q+K32E.

15 In an interesting embodiment of the invention the variant comprises, in addition to the substitutions in position 10 and 32, such as in addition to the substitutions P10Q+K32E, at least one further modification in the Gla domain.

In one preferred embodiment of the invention the further modification in the Gla domain comprises an amino acid substitution in position 33. Preferably, a hydrophobic amino
20 acid residue is introduced by substitution in position 33, such as D33I, D33L, D33M, D33V, D33F, D33Y or D33W, in particular D33F. Accordingly, in one very interesting embodiment of the invention, the variant comprises the following substitutions P10Q+K32+D33F.

In another preferred embodiment of the invention the further modification in the Gla domain comprises an insertion of at least one (such as one) amino acid residue between position
25 3 and 4. It is preferred that the inserted amino acid residue is a hydrophobic amino acid residue. Most preferably the insertion is A3AY. Accordingly, in another very interesting embodiment of the invention, the variant comprises the following modifications A3AY+P10Q+K32E or A3AY+P10Q+K32E+D33F.

In still another preferred embodiment of the invention the further modification in the
30 Gla domain comprises a substitution in position 34. It is preferred that a negatively charged amino acid residue is introduced by substitution in position 34. Most preferably the substitution is A34E. Accordingly, in still another very interesting embodiment of the invention, the variant comprises the following modifications P10Q+K32E+A34E, P10Q+K32E+D33F+A34E, A3AY+P10Q+K32E+A34E or A3AY+P10Q+K32E+D33F+A34E.

The Gla domain may also contain modifications in other positions, in particular in positions 8, 11 and 28, such as R28F or R28E. On the other hand it should be understood that the Gla domain should not be modified to such an extent that the membrane binding properties are impaired. Accordingly, it is preferred that no modifications are made in the residues that become γ -carboxylated, i.e. it is preferred that no modifications are made in residues 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35. In a similar way, it is in general not preferred that non-polypeptide moieties, such as sugar moieties and/or PEG groups, are introduced in the Gla domain. Consequently, it is preferred that no modifications are made in the Gla domain that creates an *in vivo* N-glycosylation site.

Finally, it will be understood that the modifications in the Gla domain discussed in this section must be combined with one or more of the modifications disclosed in the section entitled "*Introduction of in vivo N-glycosylation sites located outside the Gla domain*" above.

Specific examples of such "combined" variants are given below.

In one embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+T106N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A175T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+T267+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+T106N+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+T106N+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+I205T+T267N+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+T106N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A175T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+V253N.

5 In a further preferred embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+T267+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+T106N+I205T.

10 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+T106N+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+I205T+T267N+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+T106N.

15 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A175T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+I205T.

20 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+T267+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+T106N+I205T.

25 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+T106N+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+I205T+T267N+S269T.

30 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+T106N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A175T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+T267+S269T.

5 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+T106N+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+T106N+V253N.

10 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+I205T+T267N+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+T106N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+A175T.

15 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+V253N.

20 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+T267+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+T106N+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+T106N+V253N.

25 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+A34E+I205T+T267N+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E+T106N.

30 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E A175T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E T267+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E T106N+I205T.

5 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E T106N+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: P10Q+K32E+D33F+A34E I205T+T267N+S269T.

10 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E+T106N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E A175T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E I205T.

15 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E T267+S269T.

20 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E T106N+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E T106N+V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+A34E I205T+T267N+ S269T.

25 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E+T106N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E A175T.

30 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E V253N.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E T267+S269T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E T106N+I205T.

In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E T106N+V253N.

5 In a further embodiment of the invention said FVII or FVIIa variant comprises the following modifications: A3AY+P10Q+K32E+D33F+A34E I205T+T267N+ S269T.

Other modifications outside the Gla domain

In a further embodiment of the present invention, the FVII or FVIIa variant may, in
10 addition to the modifications described in the sections above, also contain mutations, which are already known to increase the intrinsic activity of the polypeptide, e.g. such as those described in WO 02/22776

Examples of preferred substitutions include substitutions selected from the group consisting of V158D, E296D, M298Q, L305V and K337A. More preferably, said substitutions
15 are selected from the group consisting of V158D+E296D+M298Q+L305V+K337A, V158D+E296D+M298Q+K337A, V158D+E296D+M298Q+L305V, V158D+E296D+M298Q, M298Q, L305V+K337A, L305V and K337A.

In a further embodiment of the present invention, the FVII or FVIIa variant may, in addition to the modifications described in the sections above, also contain mutations, which are
20 already known to cause a decreased inhibition by TFPI. One example includes the substitution K341Q disclosed by Neuenschwander et al, Biochemistry, 1995; 34:8701-8707.

Moreover, the variant may contain modifications which are believed to increase the TF binding affinity. Examples of such modifications include substitutions selected from the group consisting of L39E, L39Q, L39H, I42K, I42R, S43H, S43Q, K62E, K62R, L65Q, L65S, F71D,
25 F71Y, F71E, F71Q, F71N, E82Q, E82N, E82K and F275H

As already indicated above, the variant may also contain conservative amino acid substitutions.

The non-polypeptide moiety

30 Based on the present disclosure the skilled person will be aware that amino acid residues comprising other attachment groups may be introduced by substitution into the parent polypeptide, using the same approach as that illustrated above with *in vivo* N-glycosylation sites. For instance, one or more amino acid residues comprising an acid group (glutamic acid or

aspartic acid), tyrosine or lysine may be introduced into the positions discussed above. In particular, one or more cysteine residues may be introduced in the positions discussed above.

As indicated further above the non-polypeptide moiety of the conjugated variant is preferably selected from the group consisting of a polymer molecule, a lipophilic compound, a sugar moiety (by way of *in vivo* glycosylation) and an organic derivatizing agent. All of these agents may confer desirable properties to the variant polypeptide, in particular increased AUC_{iv}, increased functional *in vivo* half-life and/or increased plasma half-life. The variant polypeptide is normally conjugated to only one type of non-polypeptide moiety, but may also be conjugated to two or more different types of non-polypeptide moieties, e.g. to a polymer molecule and a sugar moiety, to a lipophilic group and a sugar moiety, to an organic derivatizing agent and a sugar moiety, to a lipophilic group and a polymer molecule, etc. The conjugation to two or more different non-polypeptide moieties may be done simultaneous or sequentially.

Methods of preparing a conjugated variant of the invention

In the following sections "*Conjugation to a lipophilic compound*", "*Conjugation to a polymer molecule*", "*Conjugation to a sugar moiety*" and "*Conjugation to an organic derivatizing agent*" conjugation to specific types of non-polypeptide moieties is described. In general, a conjugated variant according to the invention may be produced by culturing an appropriate host cell under conditions conducive for the expression of the variant polypeptide, and recovering the variant polypeptide, wherein a) the variant polypeptide comprises at least one N- or O-glycosylation site and the host cell is an eukaryotic host cell capable of *in vivo* glycosylation, and/or b) the variant polypeptide is subjected to conjugation to a non-polypeptide moiety *in vitro*.

It will be understood that the conjugation should be designed so as to produce the optimal molecule with respect to the number of non-polypeptide moieties attached, the size and form of such molecules (e.g. whether they are linear or branched), and the attachment site(s) in the polypeptide. The molecular weight of the non-polypeptide moiety to be used may, e.g., be chosen on the basis of the desired effect to be achieved. For instance, if the primary purpose of the conjugation is to achieve a conjugated variant having a high molecular weight (e.g. to reduce renal clearance) it is usually desirable to conjugate as few high molecular weight non-polypeptide moieties as possible to obtain the desired molecular weight. When a high degree of shielding is desirable this may be obtained by use of a sufficiently high number of low molecular weight non-polypeptide moieties (e.g. with a molecular weight of from about 300 Da to about 5 kDa, such as a molecular weight of from 300 Da to 2 kDa).

Conjugation to a polymer molecule

The polymer molecule to be coupled to the variant polypeptide may be any suitable polymer molecule, such as a natural or synthetic homo-polymer or hetero-polymer, typically with a molecular weight in the range of about 300-100,000 Da, such as about 500-20,000 Da, more preferably in the range of about 500-15,000 Da, even more preferably in the range of about 2-12 kDa, such as in the range of about 3-10 kDa. When the term "about" is used herein in connection with a certain molecular weight, the word "about" indicates an approximate average molecular weight and reflects the fact that there will normally be a certain molecular weight distribution in a given polymer preparation.

Examples of homo-polymers include a polyol (i.e. poly-OH), a polyamine (i.e. poly-NH₂) and a polycarboxylic acid (i.e. poly-COOH). A hetero-polymer is a polymer comprising different coupling groups, such as a hydroxyl group and an amine group.

Examples of suitable polymer molecules include polymer molecules selected from the group consisting of polyalkylene oxide (PAO), including polyalkylene glycol (PAG), such as polyethylene glycol (PEG) and polypropylene glycol (PPG), branched PEGs, poly-vinyl alcohol (PVA), poly-carboxylate, poly-(vinylpyrrolidone), polyethylene-co-maleic acid anhydride, polystyrene-co-maleic acid anhydride, dextran, including carboxymethyl-dextran, or any other biopolymer suitable for reducing immunogenicity and/or increasing functional *in vivo* half-life and/or serum half-life. Another example of a polymer molecule is human albumin or another abundant plasma protein. Generally, polyalkylene glycol-derived polymers are biocompatible, non-toxic, non-antigenic, non-immunogenic, have various water solubility properties, and are easily excreted from living organisms.

PEG is the preferred polymer molecule, since it has only few reactive groups capable of cross-linking compared to, e.g., polysaccharides such as dextran. In particular, mono-functional PEG, e.g. methoxypolyethylene glycol (mPEG), is of interest since its coupling chemistry is relatively simple (only one reactive group is available for conjugating with attachment groups on the polypeptide). Consequently, as the risk of cross-linking is eliminated, the resulting conjugated variants are more homogeneous and the reaction of the polymer molecules with the variant polypeptide is easier to control.

To effect covalent attachment of the polymer molecule(s) to the variant polypeptide, the hydroxyl end groups of the polymer molecule must be provided in activated form, i.e. with reactive functional groups (examples of which include primary amino groups, hydrazide (HZ), thiol, succinate (SUC), succinimidyl succinate (SS), succinimidyl succinamide (SSA),

succinimidyl propionate (SPA), succinimidyl butyrate (SBA), succinimidyl carboxymethylate (SCM), benzotriazole carbonate (BTC), N-hydroxysuccinimide (NHS), aldehyde, nitrophenylcarbonate (NPC), and tresylate (TRES)). Suitable activated polymer molecules are commercially available, e.g. from Shearwater Polymers, Inc., Huntsville, AL, USA, or from
5 PolyMASC Pharmaceuticals plc, UK.

Alternatively, the polymer molecules can be activated by conventional methods known in the art, e.g. as disclosed in WO 90/13540. Specific examples of activated linear or branched polymer molecules for use in the present invention are described in the Shearwater Polymers, Inc. 1997 and 2000 Catalogs (Functionalized Biocompatible Polymers for Research and
10 pharmaceuticals, Polyethylene Glycol and Derivatives, incorporated herein by reference).

Specific examples of activated PEG polymers include the following linear PEGs: NHS-PEG (e.g. SPA-PEG, SSPA-PEG, SBA-PEG, SS-PEG, SSA-PEG, SC-PEG, SG-PEG, and SCM-PEG), and NOR-PEG, BTC-PEG, EPOX-PEG, NCO-PEG, NPC-PEG, CDI-PEG, ALD-PEG, TRES-PEG, VS-PEG, IODO-PEG, and MAL-PEG, and branched PEGs such as
15 PEG2-NHS and those disclosed in US 5,932,462 and US 5,643,575, both of which are incorporated herein by reference. Furthermore, the following publications, incorporated herein by reference, disclose useful polymer molecules and/or PEGylation chemistries: US 5,824,778, US 5,476,653, WO 97/32607, EP 229,108, EP 402,378, US 4,902,502, US 5,281,698, US 5,122,614, US 5,219,564, WO 92/16555, WO 94/04193, WO 94/14758, WO 94/17039, WO
20 94/18247, WO 94/28024, WO 95/00162, WO 95/11924, WO95/13090, WO 95/33490, WO 96/00080, WO 97/18832, WO 98/41562, WO 98/48837, WO 99/32134, WO 99/32139, WO 99/32140, WO 96/40791, WO 98/32466, WO 95/06058, EP 439 508, WO 97/03106, WO 96/21469, WO 95/13312, EP 921 131, US 5,736,625, WO 98/05363, EP 809 996, US 5,629,384, WO 96/41813, WO 96/07670, US 5,473,034, US 5,516,673, EP 605 963, US
25 5,382,657, EP 510 356, EP 400 472, EP 183 503 and EP 154 316.

Specific examples of activated PEG polymers particularly preferred for coupling to cysteine residues, include the following linear PEGs: vinylsulfone-PEG (VS-PEG), preferably vinylsulfone-mPEG (VS-mPEG); maleimide-PEG (MAL-PEG), preferably maleimide-mPEG (MAL-mPEG) and orthopyridyl-disulfide-PEG (OPSS-PEG), preferably orthopyridyl-disulfide-
30 mPEG (OPSS-mPEG). Typically, such PEG or mPEG polymers will have a size of about 5 kDa, about 10 kD, about 12 kDa or about 20 kDa.

The conjugation of the polypeptide variant and the activated polymer molecules is conducted by use of any conventional method, e.g. as described in the following references (which also describe suitable methods for activation of polymer molecules): Harris and Zalipsky, eds.,

Poly(ethylene glycol) Chemistry and Biological Applications, AZC, Washington; R.F. Taylor, (1991), "Protein immobilisation. Fundamental and applications", Marcel Dekker, N.Y.; S.S. Wong, (1992), "Chemistry of Protein Conjugation and Crosslinking", CRC Press, Boca Raton; G.T. Hermanson et al., (1993), "Immobilized Affinity Ligand Techniques", Academic Press,
5 N.Y.).

The skilled person will be aware that the activation method and/or conjugation chemistry to be used depends on the attachment group(s) of the variant polypeptide (examples of which are given further above), as well as the functional groups of the polymer (e.g. being amine, hydroxyl, carboxyl, aldehyde, sulfhydryl, succinimidyl, maleimide, vinylsulfone or
10 haloacetate). The PEGylation may be directed towards conjugation to all available attachment groups on the variant polypeptide (i.e. such attachment groups that are exposed at the surface of the polypeptide) or may be directed towards one or more specific attachment groups, e.g. the N-terminal amino group as described in US 5,985,265 or to cysteine residues. Furthermore, the conjugation may be achieved in one step or in a stepwise manner (e.g. as described in WO
15 99/55377).

For PEGylation to cysteine residues (see above) the FVII or FVIIa variant is usually treated with a reducing agent, such as dithiothreitol (DDT) prior to PEGylation. The reducing agent is subsequently removed by any conventional method, such as by desalting. Conjugation of PEG to a cysteine residue typically takes place in a suitable buffer at pH 6-9 at temperatures
20 varying from 4°C to 25°C for periods up to 16 hours.

It will be understood that the PEGylation is designed so as to produce the optimal molecule with respect to the number of PEG molecules attached, the size and form of such molecules (e.g. whether they are linear or branched), and the attachment site(s) in the variant polypeptide. The molecular weight of the polymer to be used may e.g. be chosen on the basis of
25 the desired effect to be achieved.

In connection with conjugation to only a single attachment group on the protein (e.g. the N-terminal amino group), it may be advantageous that the polymer molecule, which may be linear or branched, has a high molecular weight, preferably about 10-25 kDa, such as about 15-25 kDa, e.g. about 20 kDa.

30 Normally, the polymer conjugation is performed under conditions aimed at reacting as many of the available polymer attachment groups with polymer molecules. This is achieved by means of a suitable molar excess of the polymer relative to the polypeptide. Typically, the molar ratios of activated polymer molecules to polypeptide are up to about 1000-1, such as up

to about 200-1, or up to about 100-1. In some cases the ration may be somewhat lower, however, such as up to about 50-1, 10-1, 5-1, 2-1 or 1-1 in order to obtain optimal reaction.

It is also contemplated according to the invention to couple the polymer molecules to the polypeptide through a linker. Suitable linkers are well known to the skilled person. A preferred example is cyanuric chloride (Abuchowski et al., (1977), J. Biol. Chem., 252, 3578-3581; US 4,179,337; Shafer et al., (1986), J. Polym. Sci. Polym. Chem. Ed., 24, 375-378).

Subsequent to the conjugation, residual activated polymer molecules are blocked according to methods known in the art, e.g. by addition of primary amine to the reaction mixture, and the resulting inactivated polymer molecules are removed by a suitable method.

It will be understood that depending on the circumstances, e.g. the amino acid sequence of the variant polypeptide, the nature of the activated PEG compound being used and the specific PEGylation conditions, including the molar ratio of PEG to polypeptide, varying degrees of PEGylation may be obtained, with a higher degree of PEGylation generally being obtained with a higher ratio of PEG to variant polypeptide. The PEGylated variant polypeptides resulting from any given PEGylation process will, however, normally comprise a stochastic distribution of conjugated polypeptide variants having slightly different degrees of PEGylation.

Coupling to a sugar moiety

In order to achieve *in vivo* glycosylation of a FVII molecule comprising one or more glycosylation sites the nucleotide sequence encoding the variant polypeptide must be inserted in a glycosylating, eucaryotic expression host. The expression host cell may be selected from fungal (filamentous fungal or yeast), insect or animal cells or from transgenic plant cells. In one embodiment the host cell is a mammalian cell, such as a CHO cell, BHK or HEK, e.g. HEK 293, cell, or an insect cell, such as an SF9 cell, or a yeast cell, e.g. *S. cerevisiae* or *Pichia pastoris*, or any of the host cells mentioned hereinafter.

Covalent *in vitro* coupling of sugar moieties (such as dextran) to amino acid residues of the variant polypeptide may also be used, e.g. as described, for example in WO 87/05330 and in Aplin et al., CRC Crit Rev. Biochem, pp. 259-306, 1981. The *in vitro* coupling of sugar moieties or PEG to protein- and peptide-bound Gln-residues can be carried out by transglutaminases (TGases). Transglutaminases catalyse the transfer of donor amine-groups to protein- and peptide-bound Gln-residues in a so-called cross-linking reaction. The donor-amine groups can be protein- or peptide-bound, such as the ϵ -amino-group in Lys-residues or it can be part of a small or large organic molecule. An example of a small organic molecule

functioning as amino-donor in TGase-catalysed cross-linking is putrescine (1,4-diaminobutane). An example of a larger organic molecule functioning as amino-donor in TGase-catalysed cross-linking is an amine-containing PEG (Sato et al., 1996, Biochemistry 35, 13072-13080).

5 TGases, in general, are highly specific enzymes, and not every Gln-residues exposed on the surface of a protein is accessible to TGase-catalysed cross-linking to amino-containing substances. On the contrary, only few Gln-residues are naturally functioning as TGase substrates but the exact parameters governing which Gln-residues are good TGase substrates remain unknown. Thus, in order to render a protein susceptible to TGase-catalysed cross-linking reactions it is often a prerequisite at convenient positions to add stretches of amino acid
10 sequence known to function very well as TGase substrates. Several amino acid sequences are known to be or to contain excellent natural TGase substrates e.g. substance P, elafin, fibrinogen, fibronectin, α_2 -plasmin inhibitor, α -caseins, and β -caseins.

Conjugation to an organic derivatizing agent

15 Covalent modification of the variant polypeptide may be performed by reacting one or more attachment groups of the variant polypeptide with an organic derivatizing agent. Suitable derivatizing agents and methods are well known in the art. For example, cysteinyl residues most commonly are reacted with α -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyl
20 residues also are derivatized by reaction with bromotrifluoroacetone, α -bromo- β -(4-imidozoyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole. Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side
25 chain. Para-bromophenacyl bromide also is useful. The reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0. Lysinyl and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyl residues. Other suitable reagents for derivatizing α -amino-containing residues include imidoesters such as methyl picolinimate, pyridoxal phosphate, pyridoxal,
30 chloroborohydride, trinitrobenzenesulfonic acid, O-methylisourea, 2,4-pentanedione and transaminase-catalyzed reaction with glyoxylate. Arginyl residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-

cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group.

Furthermore, these reagents may react with the groups of lysine as well as the arginine guanidino group. Carboxyl side groups (aspartyl or glutamyl) are selectively modified by
5 reaction with carbodiimides ($R-N=C=N-R'$), where R and R' are different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminy residues by reaction with ammonium ions.

10 *Conjugation to a lipophilic compound*

The variant polypeptide and the lipophilic compound may be conjugated to each other, either directly or by use of a linker. The lipophilic compound may be a natural compound such as a saturated or unsaturated fatty acid, a fatty acid diketone, a terpene, a prostaglandin, a
vitamine, a carotenoide or steroide, or a synthetic compound such as a carbon acid, an alcohol,
15 an amine and sulphonic acid with one or more alkyl-, aryl-, alkenyl- or other multiple unsaturated compounds. The conjugation between the variant polypeptide and the lipophilic compound, optionally through a linker may be done according to methods known in the art, e.g. as described by Bodanszky in *Peptide Synthesis*, John Wiley, New York, 1976 and in WO 96/12505.

20

Methods of preparing a polypeptide variant of the invention

The polypeptide variant of the present invention may be produced by any suitable method known in the art. Such methods include constructing a nucleotide sequence encoding the polypeptide variant and expressing the sequence in a suitable transformed or transfected
25 host. Preferably, the host cell is a gammacarboxylating host cell such as a mammalian cell. However, polypeptide variants of the invention may be produced, albeit less efficiently, by chemical synthesis or a combination of chemical synthesis or a combination of chemical synthesis and recombinant DNA technology.

A nucleotide sequence encoding a polypeptide of the invention may be constructed by
30 isolating or synthesizing a nucleotide sequence encoding the parent FVII, such as hFVII with the amino acid sequence shown in SEQ ID NO:2 and then changing the nucleotide sequence so as to effect introduction (i.e. insertion or substitution) or removal (i.e. deletion or substitution) of the relevant amino acid residue(s).

The nucleotide sequence is conveniently modified by site-directed mutagenesis in accordance with conventional methods. Alternatively, the nucleotide sequence is prepared by chemical synthesis, e.g. by using an oligonucleotide synthesizer, wherein oligonucleotides are designed based on the amino acid sequence of the desired polypeptide, and preferably selecting
5 those codons that are favored in the host cell in which the recombinant polypeptide will be produced. For example, several small oligonucleotides coding for portions of the desired polypeptide may be synthesized and assembled by PCR, ligation or ligation chain reaction (LCR) (Barany, PNAS 88:189-193, 1991). The individual oligonucleotides typically contain 5' or 3' overhangs for complementary assembly.

10 Alternative nucleotide sequence modification methods are available for producing polypeptide variants for high throughput screening, for instance methods which involve homologous cross-over such as disclosed in US 5,093,257, and methods which involve gene shuffling, i.e. recombination between two or more homologous nucleotide sequences resulting in new nucleotide sequences having a number of nucleotide alterations when compared to the
15 starting nucleotide sequences. Gene shuffling (also known as DNA shuffling) involves one or more cycles of random fragmentation and reassembly of the nucleotide sequences, followed by screening to select nucleotide sequences encoding polypeptides with desired properties. In order for homology-based nucleic acid shuffling to take place, the relevant parts of the nucleotide sequences are preferably at least 50% identical, such as at least 60% identical, more preferably
20 at least 70% identical, such as at least 80% identical. The recombination can be performed *in vitro* or *in vivo*.

Examples of suitable *in vitro* gene shuffling methods are disclosed by Stemmer et al. (1994), Proc. Natl. Acad. Sci. USA; vol. 91, pp. 10747-10751; Stemmer (1994), Nature, vol. 370, pp. 389-391; Smith (1994), Nature vol. 370, pp. 324-325; Zhao et al., Nat. Biotechnol.
25 1998, Mar; 16(3): 258-61; Zhao H. and Arnold, FB, Nucleic Acids Research, 1997, Vol. 25. No. 6 pp. 1307-1308; Shao et al., Nucleic Acids Research 1998, Jan 15; 26(2): pp. 681-83; and WO 95/17413.

An example of a suitable *in vivo* shuffling method is disclosed in WO 97/07205. Other techniques for mutagenesis of nucleic acid sequences by *in vitro* or *in vivo* recombination are
30 disclosed e.g. in WO 97/20078 and US 5,837,458. Examples of specific shuffling techniques include "family shuffling", "synthetic shuffling" and "*in silico* shuffling".

Family shuffling involves subjecting a family of homologous genes from different species to one or more cycles of shuffling and subsequent screening or selection. Family shuffling techniques are disclosed e.g. by Crameri et al. (1998), Nature, vol. 391, pp. 288-291;

Christians et al. (1999), Nature Biotechnology, vol. 17, pp. 259-264; Chang et al. (1999), Nature Biotechnology, vol. 17, pp. 793-797; and Ness et al. (1999), Nature Biotechnology, vol. 17, 893-896.

Synthetic shuffling involves providing libraries of overlapping synthetic
5 oligonucleotides based e.g. on a sequence alignment of homologous genes of interest. The synthetically generated oligonucleotides are recombined, and the resulting recombinant nucleic acid sequences are screened and if desired used for further shuffling cycles. Synthetic shuffling techniques are disclosed in WO 00/42561.

In silico shuffling refers to a DNA shuffling procedure, which is performed or
10 modelled using a computer system, thereby partly or entirely avoiding the need for physically manipulating nucleic acids. Techniques for *in silico* shuffling are disclosed in WO 00/42560.

Once assembled (by synthesis, site-directed mutagenesis or another method), the nucleotide sequence encoding the polypeptide is inserted into a recombinant vector and operably linked to control sequences necessary for expression of the FVII in the desired
15 transformed host cell.

It should of course be understood that not all vectors and expression control sequences function equally well to express the nucleotide sequence encoding the polypeptide variants described herein. Neither will all hosts function equally well with the same expression system. However, one of skill in the art may make a selection among these vectors, expression control
20 sequences and hosts without undue experimentation. For example, in selecting a vector, the host must be considered because the vector must replicate in it or be able to integrate into the chromosome. The vector's copy number, the ability to control that copy number, and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered. In selecting an expression control sequence, a variety of factors should also be
25 considered. These include, for example, the relative strength of the sequence, its controllability, and its compatibility with the nucleotide sequence encoding the polypeptide, particularly as regards potential secondary structures. Hosts should be selected by consideration of their compatibility with the chosen vector, the toxicity of the product coded for by the nucleotide sequence, their secretion characteristics, their ability to fold the polypeptide variant correctly,
30 their fermentation or culture requirements, and the ease of purification of the products coded for by the nucleotide sequence.

The recombinant vector may be an autonomously replicating vector, i.e. a vector, which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector is one which, when

introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

The vector is preferably an expression vector, in which the nucleotide sequence encoding the polypeptide variant of the invention is operably linked to additional segments
5 required for transcription of the nucleotide sequence. The vector is typically derived from plasmid or viral DNA. A number of suitable expression vectors for expression in the host cells mentioned herein are commercially available or described in the literature. Useful expression vectors for eukaryotic hosts, include, for example, vectors comprising expression control sequences from SV40, bovine papilloma virus, adenovirus and cytomegalovirus. Specific
10 vectors are, e.g., pCDNA3.1(+)\Hyg (Invitrogen, Carlsbad, CA, USA) and pCI-neo (Stratagene, La Jolla, CA, USA). Useful expression vectors for yeast cells include the 2 μ plasmid and derivatives thereof, the POT1 vector (US 4,931,373), the pJSO37 vector described in Okkels, Ann. New York Acad. Sci. 782, 202-207, 1996, and pPICZ A, B or C (Invitrogen). Useful vectors for insect cells include pVL941, pBG311 (Cate *et al.*, "Isolation of the Bovine and
15 Human Genes for Mullerian Inhibiting Substance And Expression of the Human Gene In Animal Cells", Cell, 45, pp. 685-98 (1986), pBluebac 4.5 and pMelbac (both available from Invitrogen). Useful expression vectors for bacterial hosts include known bacterial plasmids, such as plasmids from *E. coli*, including pBR322, pET3a and pET12a (both from Novagen Inc., WI, USA), wider host range plasmids, such as RP4, phage DNAs, e.g., the numerous
20 derivatives of phage lambda, e.g., NM989, and other DNA phages, such as M13 and filamentous single stranded DNA phages.

Other vectors for use in this invention include those that allow the nucleotide sequence encoding the polypeptide variant to be amplified in copy number. Such amplifiable vectors are well known in the art. They include, for example, vectors able to be amplified by DHFR
25 amplification (see, e.g., Kaufman, U.S. Pat. No. 4,470,461, Kaufman and Sharp, "Construction Of A Modular Dihydrofolate Reductase cDNA Gene: Analysis Of Signals Utilized For Efficient Expression", Mol. Cell. Biol., 2, pp. 1304-19 (1982)) and glutamine synthetase ("GS") amplification (see, e.g., US 5,122,464 and EP 338,841).

The recombinant vector may further comprise a DNA sequence enabling the vector to
30 replicate in the host cell in question. An example of such a sequence (when the host cell is a mammalian cell) is the SV40 origin of replication. When the host cell is a yeast cell, suitable sequences enabling the vector to replicate are the yeast plasmid 2 μ replication genes REP 1-3 and origin of replication.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the gene coding for dihydrofolate reductase (DHFR) or the *Schizosaccharomyces pombe* TPI gene (described by P.R. Russell, Gene 40, 1985, pp. 125-130), or one which confers resistance to a drug, e.g. ampicillin, kanamycin, tetracyclin, chloramphenicol, neomycin, hygromycin or methotrexate. For *Saccharomyces cerevisiae*, selectable markers include *ura3* and *leu2*. For filamentous fungi, selectable markers include *amdS*, *pyrG*, *arcB*, *niaD* and *sC*.

The term "control sequences" is defined herein to include all components, which are necessary or advantageous for the expression of the polypeptide variant of the invention. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide variant. Such control sequences include, but are not limited to, a leader sequence, polyadenylation sequence, propeptide sequence, promoter, enhancer or upstream activating sequence, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter.

A wide variety of expression control sequences may be used in the present invention. Such useful expression control sequences include the expression control sequences associated with structural genes of the foregoing expression vectors as well as any sequence known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof.

Examples of suitable control sequences for directing transcription in mammalian cells include the early and late promoters of SV40 and adenovirus, e.g. the adenovirus 2 major late promoter, the MT-1 (metallothionein gene) promoter, the human cytomegalovirus immediate-early gene promoter (CMV), the human elongation factor 1 α (EF-1 α) promoter, the *Drosophila* minimal heat shock protein 70 promoter, the Rous Sarcoma Virus (RSV) promoter, the human ubiquitin C (UbC) promoter, the human growth hormone terminator, SV40 or adenovirus Elb region polyadenylation signals and the Kozak consensus sequence (Kozak, M. *J Mol Biol* 1987 Aug 20;196(4):947-50).

In order to improve expression in mammalian cells a synthetic intron may be inserted in the 5' untranslated region of the nucleotide sequence encoding the polypeptide. An example of a synthetic intron is the synthetic intron from the plasmid pCI-Neo (available from Promega Corporation, WI, USA).

Examples of suitable control sequences for directing transcription in insect cells include the polyhedrin promoter, the P10 promoter, the *Autographa californica* polyhedrosis

virus basic protein promoter, the baculovirus immediate early gene 1 promoter and the baculovirus 39K delayed-early gene promoter, and the SV40 polyadenylation sequence. Examples of suitable control sequences for use in yeast host cells include the promoters of the yeast α -mating system, the yeast triose phosphate isomerase (TPI) promoter, promoters from
5 yeast glycolytic genes or alcohol dehydrogenase genes, the ADH2-4c promoter, and the inducible GAL promoter. Examples of suitable control sequences for use in filamentous fungal host cells include the ADH3 promoter and terminator, a promoter derived from the genes encoding *Aspergillus oryzae* TAKA amylase triose phosphate isomerase or alkaline protease, an *A. niger* α -amylase, *A. niger* or *A. nidulans* glucoamylase, *A. nidulans* acetamidase,
10 *Rhizomucor miehei* aspartic proteinase or lipase, the TPI1 terminator and the ADH3 terminator. Examples of suitable control sequences for use in bacterial host cells include promoters of the *lac* system, the *trp* system, the *TAC* or *TRC* system, and the major promoter regions of phage lambda.

The presence or absence of a signal peptide will, e.g., depend on the expression host
15 cell used for the production of the polypeptide variant to be expressed (whether it is an intracellular or extracellular polypeptide) and whether it is desirable to obtain secretion. For use in filamentous fungi, the signal peptide may conveniently be derived from a gene encoding an *Aspergillus* sp. amylase or glucoamylase, a gene encoding a *Rhizomucor miehei* lipase or protease or a *Humicola lanuginosa* lipase. The signal peptide is preferably derived from a gene
20 encoding *A. oryzae* TAKA amylase, *A. niger* neutral α -amylase, *A. niger* acid-stable amylase, or *A. niger* glucoamylase. For use in insect cells, the signal peptide may conveniently be derived from an insect gene (cf. WO 90/05783), such as the *Lepidopteran manduca sexta* adipokinetic hormone precursor, (cf. US 5,023,328), the honeybee melittin (Invitrogen), ecdysteroid UDPglucosyltransferase (egt) (Murphy *et al.*, Protein Expression and Purification
25 4, 349-357 (1993) or human pancreatic lipase (hpl) (Methods in Enzymology 284, pp. 262-272, 1997). A preferred signal peptide for use in mammalian cells is that of hFVII or the murine Ig kappa light chain signal peptide (Coloma, M (1992) J. Imm. Methods 152:89-104). For use in yeast cells suitable signal peptides have been found to be the α -factor signal peptide from *S. cerevisiae* (cf. US 4,870,008), a modified carboxypeptidase signal peptide (cf. L.A. Valls *et al.*,
30 Cell 48, 1987, pp. 887-897), the yeast BAR1 signal peptide (cf. WO 87/02670), the yeast aspartic protease 3 (YAP3) signal peptide (cf. M. Egel-Mitani *et al.*, Yeast 6, 1990, pp. 127-137), and the synthetic leader sequence TA57 (WO98/32867). For use in *E. coli* cells a suitable signal peptide have been found to be the signal peptide *ompA* (EP581821).

The nucleotide sequence of the invention encoding a polypeptide variant, whether prepared by site-directed mutagenesis, synthesis, PCR or other methods, may optionally include a nucleotide sequence that encode a signal peptide. The signal peptide is present when the polypeptide variant is to be secreted from the cells in which it is expressed. Such signal peptide, if present, should be one recognized by the cell chosen for expression of the polypeptide variant. The signal peptide may, e.g. be that normally associated with hFVII) or, alternatively, the signal peptide may be from another source than hFVII, such as any of those normally associated with other human wild-type vitamin K-dependent polypeptides. Furthermore, the signal peptide may be a signal peptide normally expressed from the host cell or one which is not normally expressed from the host cell. Accordingly, the signal peptide may be prokaryotic, e.g. derived from a bacterium such as *E. coli*, or eukaryotic, e.g. derived from a mammalian, or insect or yeast cell.

Any suitable host may be used to produce the polypeptide variant, including bacteria (although not particularly preferred), fungi (including yeasts), plant, insect, mammal, or other appropriate animal cells or cell lines, as well as transgenic animals or plants. Examples of bacterial host cells include grampositive bacteria such as strains of *Bacillus*, e.g. *B. brevis* or *B. subtilis*, *Pseudomonas* or *Streptomyces*, or gramnegative bacteria, such as strains of *E. coli*. The introduction of a vector into a bacterial host cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, *Molecular General Genetics* 168: 111-115), using competent cells (see, e.g., Young and Spizizin, 1961, *Journal of Bacteriology* 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, *Journal of Molecular Biology* 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, *Biotechniques* 6: 742-751), or conjugation (see, e.g., Koehler and Thorne, 1987, *Journal of Bacteriology* 169: 5771-5278). Examples of suitable filamentous fungal host cells include strains of *Aspergillus*, e.g. *A. oryzae*, *A. niger*, or *A. nidulans*, *Fusarium* or *Trichoderma*. Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* host cells are described in EP 238 023 and US 5,679,543. Suitable methods for transforming *Fusarium* species are described by Malardier *et al.*, 1989, *Gene* 78: 147-156 and WO 96/00787. Examples of suitable yeast host cells include strains of *Saccharomyces*, e.g. *S. cerevisiae*, *Schizosaccharomyces*, *Kluyveromyces*, *Pichia*, such as *P. pastoris* or *P. methanolica*, *Hansenula*, such as *H. Polymorpha* or *Yarrowia*. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, *Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology*, Volume 194, pp 182-187, Academic Press,

Inc., New York; Ito *et al.*, 1983, *Journal of Bacteriology* 153: 163; Hinnen *et al.*, 1978, *Proceedings of the National Academy of Sciences USA* 75: 1920; and as disclosed by Clontech Laboratories, Inc, Palo Alto, CA, USA (in the product protocol for the Yeastmaker™ Yeast Transformation System Kit). Examples of suitable insect host cells include a *Lepidoptera* cell
5 line, such as *Spodoptera frugiperda* (SF9 or Sf21) or *Trichoplusia ni* cells (High Five) (US 5,077,214). Transformation of insect cells and production of heterologous polypeptides therein may be performed as described by Invitrogen. Examples of suitable mammalian host cells include Chinese hamster ovary (CHO) cell lines, (e.g. CHO-K1; ATCC CCL-61), Green
10 Monkey cell lines (COS) (e.g. COS 1 (ATCC CRL-1650), COS 7 (ATCC CRL-1651)); mouse cells (e.g. NS/O), Baby Hamster Kidney (BHK) cell lines (e.g. ATCC CRL-1632 or ATCC CCL-10), and human cells (e.g. HEK 293 (ATCC CRL-1573)), as well as plant cells in tissue culture. Additional suitable cell lines are known in the art and available from public
depositories such as the American Type Culture Collection, Rockville, Maryland. Also, the mammalian cell, such as a CHO cell, may be modified to express sialyltransferase, e.g. 1,6-
15 sialyltransferase, e.g. as described in US 5,047,335, in order to provide improved glycosylation of the polypeptide variant.

In order to increase secretion it may be of particular interest to produce the polypeptide variant of the invention together with an endoprotease, in particular a PACE (paired basic amino acid converting enzyme) (e.g. as described in US 5,986,079), such as a Kex2
20 endoprotease (e.g. as described in WO 00/28065).

Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection, electroporation, DEAE-dextran mediated transfection, liposome-mediated transfection, viral vectors and the transfection method described by Life Technologies Ltd, Paisley, UK using Lipofectamin 2000. These methods are well known in the
25 art and e.g. described by Ausbel *et al.* (eds.), 1996, *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, USA. The cultivation of mammalian cells are conducted according to established methods, e.g. as disclosed in (*Animal Cell Biotechnology, Methods and Protocols*, Edited by Nigel Jenkins, 1999, Human Press Inc, Totowa, New Jersey, USA and Harrison MA and Rae IF, *General Techniques of Cell Culture*, Cambridge University Press
30 1997).

In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide variant using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory

or industrial fermenters performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared
5 according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide variant is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide variant is not secreted, it can be recovered from cell lysates.

The resulting polypeptide variant may be recovered by methods known in the art. For
10 example, the polypeptide variant may be recovered from the nutrient medium by conventional procedures including, but not limited to, centrifugation, filtration, extraction, spray drying, evaporation, or precipitation.

The polypeptides may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic,
15 chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), HPLC, or extraction (see, e.g., *Protein Purification*, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989).

Single chain polypeptide variants of the invention can be purified and activated to two-
20 chain polypeptide variants by a number of methods as described in the literature (Broze and Majerus, 1980, J. Biol. Chem. 255:1242-47 and Hedner and Kisiel, 1983, J. Clin. Invest. 71:1836-41). Another method whereby single chain polypeptide variant can be purified is by incorporation of Zn ions during purification as described in US 5,700,914. In a preferred embodiment the polypeptide variant is purified as a single chain polypeptide variant. The single
25 chain polypeptide variant is activated by either use of an immobilized enzyme (e.g. factors IIa, IXa, Xa and XIIa) or by autoactivation using a positively charged ion exchange matrix or the like.

It is advantageous to first purify the polypeptide variant in its single chain form, then PEGylate (if desired) and last activate by one of the methods described above or by
30 autoactivation as described by Pedersen et al, 1989, Biochemistry 28: 9331-36. The advantage of carrying out PEGylation before activation is that PEGylation of the new aminoterminal formed by cleavage of R152-I153 is avoided. PEGylation of this new amino terminal would render the molecule inactive since the formation of a hydrogen bond between D242 and the amino terminal of I153 is necessary for activity.

Pharmaceutical composition of the invention and its use

In a further aspect, the present invention relates to a composition, in particular to a pharmaceutical composition, comprising a polypeptide variant of the invention and a
5 pharmaceutically acceptable carrier or excipient.

The polypeptide variant or the pharmaceutical composition according to the invention may be used as a medicament.

Due to the improved properties mentioned hereinbefore, the polypeptide variants of the invention, or the pharmaceutical composition of the invention, are particular useful for the
10 treatment of uncontrollable bleeding events in trauma patients, thrombocytopenic patients, patients in anticoagulant treatment, and cirrhosis patients with variceal bleeds, or other upper gastrointestinal bleedings, and in patients undergoing orthotopic liver transplantation, or liver resection (allowing for transfusion free surgery).

Trauma is defined as an injury to living tissue caused by an extrinsic agent. It is the 4th
15 leading cause of death in the US and places a large financial burden on the economy.

Trauma is classified as either blunt or penetrative. Blunt trauma results in internal compression, organ damage and internal haemorrhage whereas penetrative trauma (as the consequence of an agent penetrating the body and destroying tissue, vessels and organs) results in external haemorrhage.

20 Haemorrhage, as a result of trauma, can start a cascade of problems. For example physiological compensation mechanisms are initiated with initial peripheral and mesenteric vasoconstriction to shunt blood to the central circulation. If circulation is not restored, hypovolemia shock (multiple organ failure due to inadequate perfusion) ensues. Since tissues throughout the body become starved for oxygen, anaerobic metabolism begins. However, the
25 concomitant lactic acid leads the blood pH to drop and metabolic acidosis develops. If acidosis is severe and uncorrected, the patient may develop multisystem failure and die.

Although the majority of trauma patients are hypothermic on arrival in the emergency room due to the environmental conditions at the scene, inadequate protection, intravenous fluid administration and ongoing blood loss worsen the hypothermic state. Deficiencies in
30 coagulation factors can result from blood loss or transfusions. Meanwhile, acidosis and hypothermia interfere with blood clotting mechanisms. Thus coagulopathy develops, which in turn, may mask surgical bleeding sites and hamper the control of mechanical bleeding. Hypothermia, coagulopathy and acidosis are often characterised as the "trauma triad of death"

Trauma may be caused by several events. For example, road traffic accidents result in many different types of trauma. Whilst some road traffic accidents are likely to result in penetrative trauma, many road traffic accidents are likely to inflict blunt trauma to both head and body. However, these various types of trauma can all result in coagulopathy in the patient.

5 Road traffic accidents are the leading cause of accidental death in the US. There are over 42,000 deaths from them in the US each year. Many trauma patients die at the location of the accident either whilst being treated by the paramedics, before they arrive or in transit to the ER.

Another example includes gunshot wounds. Gunshot wounds are traumas that can result in massive bleeding. They are penetrative and destroy tissue as the bullet passes through
10 the body, whether it be in the torso or a limb. In the US about 40,000 people a year die from gunshot wounds

A further example includes falls. Falls result in a similar profile of trauma type to road traffic accidents. By falling onto a solid object or the ground from height can cause both penetrative and decelerative blunt trauma. In the US, falls are a common cause of accidental
15 death, numbering about 13,000.

A still further example includes machinery accidents. A smaller number of people die in the US from machinery accident related deaths, whether struck by, or entangled in machinery. The figures are small but significant – around 2,000.

A still further example includes stab wounds. Stab wounds are penetrative injuries that
20 can also cause massive bleeding. The organs most likely to be damaged in a stab wound are the liver, small intestine and the colon.

Cirrhosis of the liver is the terminal sequel of prolonged repeated injury to the hepatic parenchyma. The end result is the formation of broad bands of fibrous tissue separating regenerative nodules that do not maintain the normal organization of liver lobules and thus
25 cause deteriorated liver function. Patients have prolonged prothrombin times as a result of the depletion of vitamin K-dependent coagulation factors. Pathogenetically, liver cirrhosis should be regarded as the final common pathway of chronic liver injury, which can result from any form of intense repeated prolonged liver cell injury. Cirrhosis of the liver may be caused by direct liver injury, including chronic alcoholism, chronic viral hepatitis (types B, C, and D), and
30 auto immune hepatitis as well as by indirect injury by way of bile duct damage, including primary biliary cirrhosis, primary sclerosing cholangitis and biliary atresia. Less common causes of cirrhosis include direct liver injury from inherited disease such as cystic fibrosis, alpha-1-antitrypsin deficiency, hemochromatosis, Wilson's disease, galactosemia, and glycogen storage disease.

Transplantation is primarily reserved for late stage cirrhotic patients, where it is the key intervention for treating the disease. To be eligible for transplantation, a patient must be classified as Child's B or C, as well as meet additional criteria for selection. Last year, in the US alone, 4,954 transplants were performed.

5 It has been estimated that there are 6,000 bleeding episodes associated with patients undergoing resection each year. This correlates with the reserved position of this procedure although seems slightly high in comparison with transplantation numbers.

Accurate data on the incidence of variceal bleeding is hard to obtain. The key facts known are that at the time of diagnosis, varices are present in about 60% of decompensated and
10 30% of compensated patients and that about 30% of these patients with varices will experience a bleed and that each episode of variceal bleeding is associated with a 30% risk of mortality.

Thus, in a further aspect the present invention relates to a polypeptide variant of the invention for the manufacture of a medicament for the treatment of diseases or disorder wherein clot formation is desirable. A still further aspect of the present invention relates to a method for
15 treating a mammal having a disease or disorder wherein clot formation is desirable, comprising administering to a mammal in need thereof an effective amount of the polypeptide variant or the pharmaceutical composition of the invention.

Thrombocytopenia is caused by one of three mechanisms-decreased bone marrow production, increased splenic sequestration, or accelerated destruction of platelets.

20 Thrombocytopenia is a risk factor for hemorrhage, and platelet transfusion reduces the incidence of bleeding. The threshold for prophylactic platelet transfusion is 10,000/ μ l. In patients without fever or infections, a threshold of 5000/ μ l may be sufficient to prevent spontaneous hemorrhage. For invasive procedures, 50,000/ μ l platelets is the usual target level. In patients who develop antibodies to platelets following repeated transfusions, bleeding can be
25 extremely difficult to control.

Examples of diseases/disorders wherein increased clot formation is desirable include, but is not limited to, hemorrhages, including brain hemorrhages, as well as patient with severe uncontrolled bleedings, such as trauma. Further examples include patients undergoing living transplantations, patients undergoing resection and patients with variceal bleedings.

30 The polypeptide variants of the invention is administered to patients in a therapeutically effective dose, normally one approximately paralleling that employed in therapy with rFVII such as NovoSeven®, or at lower dosage. By "therapeutically effective dose" herein is meant a dose that is sufficient to produce the desired effects in relation to the condition for

which it is administered. The exact dose will depend on the circumstances, and will be ascertainable by one skilled in the art using known techniques. Normally, the dose should be capable of preventing or lessening the severity or spread of the condition or indication being treated. It will be apparent to those of skill in the art that an effective amount of a polypeptide variant or composition of the invention depends, *inter alia*, upon the disease, the dose, the administration schedule, whether the polypeptide variant or composition is administered alone or in conjunction with other therapeutic agents, the plasma half-life of the compositions, and the general health of the patient. Preferably, the polypeptide variant or composition of the invention is administered in an effective dose, in particular a dose which is sufficient to normalize the coagulation disorder.

The polypeptide variant of the invention is preferably administered in a composition including a pharmaceutically acceptable carrier or excipient. "Pharmaceutically acceptable" means a carrier or excipient that does not cause any untoward effects in patients to whom it is administered. Such pharmaceutically acceptable carriers and excipients are well known in the art (see, for example, Remington's Pharmaceutical Sciences, 18th edition, A. R. Gennaro, Ed., Mack Publishing Company [1990]; Pharmaceutical Formulation Development of Peptides and Proteins, S. Frokjaer and L. Hovgaard, Eds., Taylor & Francis [2000] ; and Handbook of Pharmaceutical Excipients, 3rd edition, A. Kibbe, Ed., Pharmaceutical Press [2000]).

The polypeptide variant of the invention can be formulated into pharmaceutical compositions by well-known methods. Suitable formulations are described by Remington's Pharmaceutical Sciences by E.W. Martin (Mark Publ. Co., 16th Ed., 1980).

The polypeptide variant of the invention can be used "as is" and/or in a salt form thereof. Suitable salts include, but are not limited to, salts with alkali metals or alkaline earth metals, such as sodium, potassium, calcium and magnesium, as well as e.g. zinc salts. These salts or complexes may be present as a crystalline and/or amorphous structure.

The pharmaceutical composition of the invention may be administered alone or in conjunction with other therapeutic agents. These agents may be incorporated as part of the same pharmaceutical composition or may be administered separately from the polypeptide variant of the invention, either concurrently or in accordance with another treatment schedule. In addition, the polypeptide variant or pharmaceutical composition of the invention may be used as an adjuvant to other therapies.

A "patient" for the purposes of the present invention includes both humans and other mammals. Thus, the methods are applicable to both human therapy and veterinary applications.

The pharmaceutical composition comprising the polypeptide variant of the invention may be formulated in a variety of forms, e.g. as a liquid, gel, lyophilized, or as a compressed solid. The preferred form will depend upon the particular indication being treated and will be apparent to one skilled in the art.

5 In particular, the pharmaceutical composition comprising the polypeptide variant of the invention may be formulated in lyophilised or stable soluble form. The polypeptide variant may be lyophilised by a variety of procedures known in the art. The polypeptide variant may be in a stable soluble form by the removal or shielding of proteolytic degradation sites as described herein. The advantage of obtaining a stable soluble preparation lies in easier handling for the
10 patient and, in the case of emergencies, quicker action, which potentially can become life saving. The preferred form will depend upon the particular indication being treated and will be apparent to one of skill in the art.

The administration of the formulations of the present invention can be performed in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously,
15 intracerebrally, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, intraocularly, or in any other acceptable manner. The formulations can be administered continuously by infusion, although bolus injection is acceptable, using techniques well known in the art, such as pumps or implantation. In some instances the formulations may be directly applied as a solution or spray.

20

Parentals

A preferred example of a pharmaceutical composition is a solution designed for parenteral administration. Although in many cases pharmaceutical solution formulations are provided in liquid form, appropriate for immediate use, such parenteral formulations may also
25 be provided in frozen or in lyophilized form. In the former case, the composition must be thawed prior to use. The latter form is often used to enhance the stability of the active compound contained in the composition under a wider variety of storage conditions, as it is recognized by those skilled in the art that lyophilized preparations are generally more stable than their liquid counterparts. Such lyophilized preparations are reconstituted prior to use by the
30 addition of one or more suitable pharmaceutically acceptable diluents such as sterile water for injection or sterile physiological saline solution.

In case of parenterals, they are prepared for storage as lyophilized formulations or aqueous solutions by mixing, as appropriate, the polypeptide variant having the desired degree of purity with one or more pharmaceutically acceptable carriers, excipients or stabilizers

typically employed in the art (all of which are termed "excipients"), for example buffering agents, stabilizing agents, preservatives, isotoniifiers, non-ionic surfactants or detergents, antioxidants and/or other miscellaneous additives.

Buffering agents help to maintain the pH in the range which approximates physiological conditions. They are typically present at a concentration ranging from about 2 mM to about 50 mM. Suitable buffering agents for use in the present invention include both organic and inorganic acids and salts thereof such as citrate buffers (e.g., monosodium citrate-disodium citrate mixture, citric acid-trisodium citrate mixture, citric acid-monosodium citrate mixture, etc.), succinate buffers (e.g., succinic acid-monosodium succinate mixture, succinic acid-sodium hydroxide mixture, succinic acid-disodium succinate mixture, etc.), tartrate buffers (e.g., tartaric acid-sodium tartrate mixture, tartaric acid-potassium tartrate mixture, tartaric acid-sodium hydroxide mixture, etc.), fumarate buffers (e.g., fumaric acid-monosodium fumarate mixture, fumaric acid-disodium fumarate mixture, monosodium fumarate-disodium fumarate mixture, etc.), gluconate buffers (e.g., gluconic acid-sodium glyconate mixture, gluconic acid-sodium hydroxide mixture, gluconic acid-potassium glyconate mixture, etc.), oxalate buffer (e.g., oxalic acid-sodium oxalate mixture, oxalic acid-sodium hydroxide mixture, oxalic acid-potassium oxalate mixture, etc.), lactate buffers (e.g., lactic acid-sodium lactate mixture, lactic acid-sodium hydroxide mixture, lactic acid-potassium lactate mixture, etc.) and acetate buffers (e.g., acetic acid-sodium acetate mixture, acetic acid-sodium hydroxide mixture, etc.). Additional possibilities are phosphate buffers, histidine buffers and trimethylamine salts such as Tris.

Stabilizers refer to a broad category of excipients, which can range in function from a bulking agent to an additive which solubilizes the therapeutic agent or helps to prevent denaturation or adherence to the container wall. Typical stabilizers can be polyhydric sugar alcohols (enumerated above); amino acids such as arginine, lysine, glycine, glutamine, asparagine, histidine, alanine, ornithine, L-leucine, 2-phenylalanine, glutamic acid, threonine, etc., organic sugars or sugar alcohols, such as lactose, trehalose, stachyose, mannitol, sorbitol, xylitol, ribitol, myoinositol, galactitol, glycerol and the like, including cyclitols such as inositol; polyethylene glycol; amino acid polymers; sulfur-containing reducing agents, such as urea, glutathione, thioctic acid, sodium thioglycolate, thioglycerol, α -monothioglycerol and sodium thiosulfate; low molecular weight polypeptides (i.e. <10 residues); proteins such as human serum albumin, bovine serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; monosaccharides such as xylose, mannose, fructose and glucose; disaccharides such as lactose, maltose and sucrose; trisaccharides such as raffinose, and

polysaccharides such as dextran. Stabilizers are typically present in the range of from 0.1 to 10,000 parts by weight based on the active protein weight.

Preservatives are added to retard microbial growth, and are typically added in amounts of about 0.2%-1% (w/v). Suitable preservatives for use with the present invention include
5 phenol, benzyl alcohol, meta-cresol, methyl paraben, propyl paraben, octadecyldimethylbenzyl ammonium chloride, benzalkonium halides (e.g. benzalkonium chloride, bromide or iodide), hexamethonium chloride, alkyl parabens such as methyl or propyl paraben, catechol, resorcinol, cyclohexanol and 3-pentanol.

Isotonicifiers are added to ensure isotonicity of liquid compositions and include
10 polyhydric sugar alcohols, preferably trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol and mannitol. Polyhydric alcohols can be present in an amount between 0.1% and 25% by weight, typically 1% to 5%, taking into account the relative amounts of the other ingredients.

Non-ionic surfactants or detergents (also known as "wetting agents") may be present to
15 help solubilizing the therapeutic agent as well as to protect the therapeutic polypeptide against agitation-induced aggregation, which also permits the formulation to be exposed to shear surface stress without causing denaturation of the polypeptide. Suitable non-ionic surfactants include polysorbates (20, 80, etc.), polyoxamers (184, 188 etc.), Pluronic® polyols, polyoxyethylene sorbitan monoethers (Tween®-20, Tween®-80, etc.).

20 Additional miscellaneous excipients include bulking agents or fillers (e.g. starch), chelating agents (e.g. EDTA), antioxidants (e.g., ascorbic acid, methionine, vitamin E) and cosolvents.

The active ingredient may also be entrapped in microcapsules prepared, for example, by coascervation techniques or by interfacial polymerization, for example
25 hydroxymethylcellulose, gelatin or poly-(methylmethacrylate) microcapsules, in colloidal drug delivery systems (for example liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, *supra*.

Parenteral formulations to be used for *in vivo* administration must be sterile. This is
30 readily accomplished, for example, by filtration through sterile filtration membranes.

Sustained release preparations

Examples of sustained-release preparations include semi-permeable matrices of solid hydrophobic polymers containing the polypeptide variant, the matrices having a suitable form such as a film or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate) or poly(vinylalcohol)), polylactides, copolymers of L-glutamic acid and ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the ProLease® technology or Lupron Depot® (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for long periods such as up to or over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated polypeptides remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37°C, resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

The invention is further described in the following non-limiting examples.

MATERIALS AND METHODS*Accessible Surface Area (ASA)*

The computer program Access (B. Lee and F.M.Richards, J. Mol.Biol. 55: 379-400 (1971)) version 2 (© 1983 Yale University) is used to compute the accessible surface area (ASA) of the individual atoms in the structure. This method typically uses a probe-size of 1.4Å and defines the Accessible Surface Area (ASA) as the area formed by the center of the probe. Prior to this calculation all water molecules and all hydrogen atoms should be removed from the coordinate set, as should other atoms not directly related to the protein.

Fractional ASA of Side Chain

The fractional ASA of the side chain atoms is computed by division of the sum of the ASA of the atoms in the side chain with a value representing the ASA of the side chain atoms of that residue type in an extended Ala-x-Ala tripeptide (See Hubbard, Campbell & Thornton
 5 (1991) J.Mol.Biol.220,507-530). For this example the CA atom is regarded as a part of the side chain of Glycine residues but not for the remaining residues. The following table is used as standard 100% ASA for the side chain:

Ala	69.23 Å ²		Leu	140.76 Å ²
Arg	200.35 Å ²		Lys	162.50 Å ²
Asn	106.25 Å ²		Met	156.08 Å ²
Asp	102.06 Å ²		Phe	163.90 Å ²
Cys	96.69 Å ²		Pro	119.65 Å ²
Gln	140.58 Å ²		Ser	78.16 Å ²
Glu	134.61 Å ²		Thr	101.67 Å ²
Gly	32.28 Å ²		Trp	210.89 Å ²
His	147.00 Å ²		Tyr	176.61 Å ²
Ile	137.91 Å ²		Val	114.14 Å ²

10 Residues not detected in the structure are defined as having 100% exposure as they are thought to reside in flexible regions. The gamma-carboxy glutamic acids at positions 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35 are all defined as being 100% exposed.

Determining Distances Between Atoms

15 The distance between atoms is most easily determined using molecular graphics software e.g. InsightII® v. 98.0, MSI INC.

Active Site Region

20 The active site region is defined as any residues having at least one atom within 10 Å of any atom in the catalytic triad (residues H193, D242, S344).

Determination of Tissue Factor Binding Site

The TF binding site is defined as comprising of all residues having their accessible surface area changed upon TF binding. This is determined by at least two ASA calculations; one on the isolated ligand(s) in the ligand(s)/receptor(s) complex and one on the complete
5 ligand(s)/receptor(s) complex.

Measurement of Reduced Sensitivity to Proteolytic Degradation

Proteolytic degradation can be measured using the assay described in US 5,580,560, Example 5, where proteolysis is autoprolysis.

10 Furthermore, reduced proteolysis can be tested in an *in vivo* model using radiolabelled samples and comparing proteolysis of rhFVIIa and the polypeptide variant of the invention by withdrawing blood samples and subjecting these to SDS-PAGE and autoradiography.

Irrespective of the assay used for determining proteolytic degradation, "reduced proteolytic degradation" is intended to mean a measurable reduction in cleavage compared to
15 that obtained by rhFVIIa as measured by gel scanning of Coomassie stained SDS-PAGE gels, HPLC or as measured by conserved catalytic activity in comparison to wild type using the tissue factor independent activity assay described below.

Determination of the Molecular Weight of Polypeptide Variants

20 The molecular weight of polypeptide variants is determined by either SDS-PAGE, gel filtration, Western Blots, matrix assisted laser desorption mass spectrometry or equilibrium centrifugation, e.g. SDS-PAGE according to Laemmli, U.K., Nature Vol 227 (1970), pp. 680-85.

TF-independent Factor X Activation Assay

25 This assay has been described in detail on page 39826 in Nelsestuen et al., J Biol Chem, 2001; 276:39825-39831.

Briefly, the molecule to be assayed (either hFVIIa, rhFVIIa or the polypeptide variant of the invention in its activated form) is mixed with a source of phospholipid (phosphatidylcholine and phosphatidylserine in a ratio of 8:2 or phosphatidylcholine, phosphatidylserine and
30 phosphatidylethanol in a ratio of 4:2:4) and Factor X in Tris buffer containing BSA. After a specified incubation time the reaction is stopped by addition of excess EDTA. The concentration of factor Xa is then measured from absorbance change at 405 nm after addition of a chromogenic substrate (S-2222, Chromogenix). After correction from background the tissue factor independent activity of rhFVIIa (a_{wt}) is determined as the absorbance change after 10 minutes and the tissue

factor independent activity of the polypeptide variant of the invention (a_{variant}) is also determined as the absorbance change after 10 minutes. The ratio between the activity of the polypeptide variant, in its activated form, and the activity of rhFVIIa is defined as $a_{\text{variant}}/a_{\text{wt}}$.

5 *Clotting Assay*

Clotting activity is measured in one-stage assays and clotting times are recorded on a Thrombotrack IV coagulometer (MEDINOR). FVII depleted human plasma (American Diagnostica) is reconstituted and equilibrated at room temperature for 15-20 minutes. 50 μ l of plasma is then transferred to the coagulometer cups.

10 hFVIIa, rhFVIIa or variants are diluted in Glyoxaline Buffer (5.7 mM barbiturate, 4.3 mM sodium citrate, 117 mM NaCl, 1 mg/mL BSA, pH 7.35). The samples are added to the cup in 50 μ l and incubated at 37°C for 2 minutes.

Thromboplastin (MEDINOR) is reconstituted with water and CaCl_2 is added. The reaction is initiated by adding 0.1 ml thromboplastin containing 4.5 mM CaCl_2 .

15 Data are analysed using PRISM software.

TF-independent Clotting Assay

This assay is performed as described above under "*Clotting Assay*" but without addition of thromboplastin.

20

Amidolytic Assay

The ability of the variants to cleave small peptide substrates can be measured using the chromogenic substrate S-2288 (D-Ile-Pro-Arg-p-nitroanilide). The amidolytic activity may be measured both in the presence and absence of antithrombin III (ATIII).

25 HFVIIa, rhFVIIa or variant is diluted to 90 nM in assay buffer (50 mM Na-Hepes pH 7.5, 150 mM NaCl, 5 mM CaCl_2 , 0.1% BSA, 1U/ml Heparin). Furthermore, soluble TF (sTF) is diluted to 450 nM in assay buffer. ATIII is diluted to 900 nM in assay buffer. 120 μ l of assay buffer is mixed with 20 μ l of the FVIIa sample, 20 μ l sTF and 20 μ l ATIII or assay buffer. The final concentrations of FVIIa, sTF and ATIII are 10, 50 and 100 nM, respectively. After 5 min
30 incubation at room temperature with gentle shaking, followed by 10 min incubation at 37°C, the reaction is started by addition of the S-2288 substrate to 1 mM and the absorption at 405 nm is determined at several time points.

Thrombogram Assay

The effect of hFVIIa, rhFVIIa or variant on thrombin generation in human plasma is tested in a modified version of the assay described on page 589 in Hemker et al. in *Thromb Haemost* 2000; 83: 589-91. Briefly, the molecule to be assayed (either hFVIIa, rhFVIIa or
5 variant) is mixed with normal platelet rich plasma (PRP), normal platelet poor plasma (PPP) or FVII depleted PPP with or without the addition of recombinant human tissue factor (rTF), relipidated rTF or another source of TF (such as thromboplastin). A source of phospholipid (phosphatidylcholine and phosphatidylserine in a ratio of 8:2 or phosphatidylcholine, phosphatidylserine and phosphatidylethanol in a ratio of 4:2:4) can be added.

10 The reaction is started by addition of a fluoregenic thrombin substrate and calcium chloride. The fluorescence is measured continuously and the thrombin amidolytic activity is calculated by calculating the slope of the fluorescence curve (the increase in fluorescence over time). In this way the time until maximum thrombin amidolytic activity is obtained (T_{max}) and total thrombin work (area under the curve (AUC)) can be calculated.

15 The following procedure is used: PRP is obtained by centrifuging freshly drawn blood at 250 g, 15°C for 10 min. Blood coagulation is inhibited either by using citrate (13 mM tri-sodium citrate), corn trypsin inhibitor (50-100 µg/ml blood) or a combination of citrate and corn trypsin inhibitor. The platelet count is adjusted to 3×10^8 /mL using buffer or autologous platelet poor plasma (PPP). PPP is obtained by double centrifugation of PRP at 1000 g, 15°C
20 for 10 min. FVII depletion is done by incubating PPP with a FVII specific monoclonal antibody coupled to a solid phase.

Per well of a 96-well microtiter plate 80 µl PRP is added and 20µL buffer containing rhFVII or variant to be tested in final concentrations between 0.1 and 100 nM. rTF is added in 5 µL assay buffer to a final concentration of 1 pM. The assay buffer consists of 20mM Hepes,
25 150mM NaCl and 60mg/ml BSA in distilled water. The reaction is started by adding 20µL of the substrate solution containing 0.1 M calcium chloride. The assay plate and reagents are prewarmed to 37°C and the reaction takes place at this temperature. The fluorimeter used is a BMG Fluormeter with a excitation filter at 390 nm and an emission filter at 460 nm. The fluorescence is measured in each well of 96-well clear bottom plates in 20-40 second interval
30 over 30-180 minutes. Data are analyzed using PRISM Software.

ELISA Assay

FVII/FVIIa (or variant) concentrations are determined by ELISA. Wells of a microtiter plate are coated with an antibody directed against the protease domain using a solution of 2 µg/ml in PBS (100 µl per well). After 2 hours coating at R.T., the wells are washed 4 times
5 with THT buffer (100 mM NaCl, 50 mM Tris-HCl pH 7.2 0.05% Tween-20). Subsequently, 200 µl of 1% Casein (diluted from 2.5% stock using 100 mM NaCl, 50 mM Tris-HCl pH 7.2) is added per well for blocking. After 1 hr incubation at R.T., the wells are emptied, and 100 µl of sample (optionally diluted in dilution buffer (THT + 0.1% Casein)) is added. After another incubation of 1 hr at room temperature, the wells are washed 4 times with THT buffer, and 100
10 µl of a biotin-labelled antibody directed against the EGF-like domain (1 µg/ml) is added. After another 1 hr incubation at R.T., followed by 4 more washes with THT buffer, 100 µl of streptavidin-horse radish peroxidase (DAKO A/S, Glostrup, Denmark, 1/10000 diluted) is added. After another 1 hr incubation at R.T., followed by 4 more washes with THT buffer, 100 µl of TMB (3,3',5,5'-tetramethylbenzidine, Kem-en-Tech A/S, Denmark) is added. After 30
15 min incubation at R.T. in the dark, 100 µl of 1 M H₂SO₄ is added and OD_{450nm} is determined. A standard curve is prepared using rhFVIIa (NovoSeven®).

Alternatively, FVII/FVIIa or variants may be quantified through the Gla domain rather than through the protease domain. In this ELISA set-up, wells are coated overnight with an antibody directed against the EGF-like domain and for detection, a calcium-dependent biotin-
20 labelled monoclonal anti-Gla domain antibody is used (2 µg/ml, 100 µl per well). In this set-up, 5 mM CaCl₂ is added to the THT and dilution buffers.

Whole Blood Assay

The whole blood clotting assay is performed as described by Elg et al. Thrombosis
25 Res. 2001, 101(3):159-170.

Reconstituted Coagulation Assay

The reconstituted coagulation assay is performed as described by Van't Veer et al. Blood 2000, 95(4), 1330-1335.

EXAMPLES**Example 1**

The X-ray structure of hFVIIa in complex with soluble tissue factor by Banner et.al., J
 5 Mol Biol, 1996; 285:2089 is used for this example. It is noted that the numbering of residues in
 the reference does not follow the sequence. Here we have used the sequential numbering
 according to SEQ ID NO:2. The gamma-carboxy glutamic acids at positions 6, 7, 14, 16, 19,
 20, 25, 26, 29 and 35 are all here named GLU (three letter abbreviation) or E (one letter
 abbreviation). Residues 143-152 are not present in the structure.

10

Surface Exposure

Performing fractional ASA calculations on FVII fragments alone combined with the
 definition of accessibilities of non standard and/or missing residues described in the methods
 resulted in the following residues having more than 25% of their side chain exposed to the
 15 surface: A1, N2, A3, F4, L5, E6, E7, L8, R9, P10, S12, L13, E14, E16, K18, E19, E20, Q21,
 S23, F24, E25, E26, R28, E29, F31, K32, D33, A34, E35, R36, K38, L39, W41, I42, S43, S45,
 G47, D48, Q49, A51, S52, S53, Q56, G58, S60, K62, D63, Q64, L65, Q66, S67, I69, F71, L73,
 P74, A75, E77, G78, R79, E82, T83, H84, K85, D86, D87, Q88, L89, I90, V92, N93, E94,
 G97, E99, S103, D104, H105, T106, G107, T108, K109, S111, R113, E116, G117, S119, L120,
 20 L121, A122, D123, G124, V125, S126, T128, P129, T130, V131, E132, I140, L141, E142,
 K143, R144, N145, A146, S147, K148, P149, Q150, G151, R152, G155, K157, V158, P160,
 K161, E163, L171, N173, G174, A175, N184, T185, I186, H193, K197, K199, N200, R202,
 N203, I205, S214, E215, H216, D217, G218, D219, S222, R224, S232, T233, V235, P236,
 G237, T238, T239, N240, H249, Q250, P251, V253, T255, D256, E265, R266, T267, E270,
 25 R271, F275, V276, R277, F278, L280, L287, L288, D289, R290, G291, A292, T293, L295,
 E296, N301, M306, T307, Q308, D309, L311, Q312, Q313, R315, K316, V317, G318, D319,
 S320, P321, N322, T324, E325, Y326, Y332, S333, D334, S336, K337, K341, G342, H351,
 R353, G354, Q366, G367, T370, V371, G372, R379, E385, Q388, K389, R392, S393, E394,
 P395, R396, P397, G398, V399, L400, L401, R402, P404 and P406 (A1-S45 are located in the
 30 Gla domain, the remaining positions are located outside the Gla domain).

The following residues had more than 50% of their side chain exposed to the surface:
 A1, A3, F4, L5, E6, E7, L8, R9, P10, E14, E16, K18, E19, E20, Q21, S23, E25, E26, E29, K32,
 A34, E35, R36, K38, L39, I42, S43, G47, D48, A51, S52, S53, Q56, G58, S60, K62, L65, Q66,
 S67, I69, F71, L73, P74, A75, E77, G78, R79, E82, H84, K85, D86, D87, Q88, L89, I90, V92,

N93, E94, G97, T106, G107, T108, K109, S111, E116, S119, L121, A122, D123, G124, V131, E132, L141, E142, K143, R144, N145, A146, S147, K148, P149, Q150, G151, R152, G155, K157, P160, N173, G174, A175, K197, K199, N200, R202, S214, E215, H216, G218, R224, V235, P236, G237, T238, H249, Q250, V253, D256, T267, F275, R277, F278, L288, D289,
 5 R290, G291, A292, T293, L295, N301, M306, Q308, D309, L311, Q312, Q313, R315, K316, G318, D319, N322, E325, D334, K341, G354, G367, V371, E385, K389, R392, E394, R396, P397, G398, R402, P404 and P406 (A1-S43 are located in the Gla domain, the remaining positions are located outside the Gla domain)..

10 *Tissue Factor Binding Site*

Performing ASA calculations the following residues in human FVII change their ASA in the complex. These residues were defined as constituting the receptor binding site: L13, K18, F31, E35, R36, L39, F40, I42, S43, S60, K62, D63, Q64, L65, I69, C70, F71, C72, L73, P74, F76, E77, G78, R79, E82, K85, Q88, I90, V92, N93, E94, R271, A274, F275, V276, R277,
 15 F278, R304, L305, M306, T307, Q308, D309, Q312, Q313, E325 and R379.

Active Site Region

The active site region is defined as any residue having at least one atom within a distance of 10 Å from any atom in the catalytic triad (residues H193, D242, S344): I153, Q167,
 20 V168, L169, L170, L171, Q176, L177, C178, G179, G180, T181, V188, V189, S190, A191, A192, H193, C194, F195, D196, K197, I198, W201, V228, I229, I230, P231, S232, T233, Y234, V235, P236, G237, T238, T239, N240, H241, D242, I243, A244, L245, L246, V281, S282, G283, W284, G285, Q286, T293, T324, E325, Y326, M327, F328, D338, S339, C340, K341, G342, D343, S344, G345, G346, P347, H348, L358, T359, G360, I361, V362, S363,
 25 W364, G365, C368, V376, Y377, T378, R379, V380, Q382, Y383, W386, L387, L400 and F405.

The Ridge of the Active Site Binding Cleft

The ridge of the active site binding cleft region was defined by visual inspection of the
 30 FVIIa structure 1FAK.pdb as: N173, A175, K199, N200, N203, D289, R290, G291, A292, P321 and T370.

Example 2*Design of an expression cassette for expression of rhFVII in mammalian cells*

The DNA sequence shown in SEQ ID NO:1, encompassing the short form of the full length cDNA encoding hFVII with its native short signal peptide (Hagen et al., 1986. PNAS 83:2412), was synthesized in order to facilitate high expression in mammalian cells. First the ATG start codon context was modified according to the Kozak consensus sequence (Kozak, M. *J Mol Biol* 1987 Aug 20;196(4):947-50), so that there is a perfect match to the consensus sequence upstream of the ATG start codon. Secondly the open reading frame of the native cDNA was modified by making a bias in the codon usage towards the codons frequently used in highly expressed human genes. Further, two translational stop codons were inserted at the end of the open reading frame in order to facilitate efficient translational stop. The fully synthetic and expression optimized hFVII gene was assembled from 70-mer DNA oligonucleotides and finally amplified using end primers inserting *Bam*HI and *Hind*III sites at the 5' and 3' ends respectively using standard PCR techniques, which resulted in the following sequence:

```

15 ggatccgccaccatggtcagccaggccctccgcctcctgtgcctgctcctggggctgcagggtgcctggctgccgtcttcgtcaccca
   ggaggaagccatggcgtcctgcacgccggcgccgggccaatgcctttctggaagagctccgccctggctccctggaacgcgaatgc
   aaagaggaacagtgcagcttgaggaagcccgggagatttcaagacgctgagcggaccaaactgtttggattagctatagcgaatggc
   gatcagtgccctccagccctgccagaacgggggctcctgcaaagaccagctgcagagctatatctgcttctgcctgcctgctttagg
   ggcgcaattgcgaaccataaggtatgaccagctgattgcgtcaacgaaaacgggggctgcgagcagtactgcagcgatcacacggg
20 cacgaagcggagctgccgtgccacgaaggctatagcctcctggctgacggggtgctcgcacgccacgggtggaatacccttgcggg
   aagattcccatttagaaaagcggaacgctagcaaaccacgggcccggatcgctggcggaaggctgcctaaaggaggagtgcccct
   ggcaggtcctgctcctggtcaacggggccagctgtgcggcgggaccctcatcaatacatttgggtcgtgtccgcgctcactgcttcg
   ataagattaagaattggcggaacctcatcgctgtcgcggaacacgatctgctcgagcatgacggggacgaacagctcccgccgggtg
   gctcaggtcatcattccctccacctatgtgcctggcagcaaatcacgatatgctctgctccgctccaccagcccgtcgtgctcaccga
25 tcacgtcgtgcctctgtgcctgcctgagcggaccttagcgaacgcacgctggctttcgtccgcttagcctcgttccggctggggccag
   ctgctcgaccggggcgctaccgctctcgagctgatggtgctcaacgtccccggctgatgaccaggactgcctgcagcagctcccgcaa
   agtgggggactcccccaatatcacggagtatatgtttgcgctggctatagcgaatggtccaaggatagctgcaagggggactccggcg
   ggccccatgccacgcactatcggggacctgtacctcacgggacgtcagctggggccagggtgcgccacgggtggggcactttg
   gcgtctacacgcgcgtcagccagtacattgagtggtgcagaagctcatgcggagcgaaccccgcccggggtgctcctgcggggccc
30 cttcccttgataaaagctt

```

A vector for the cloning of the generated PCR product encompassing the expression cassette for hFVII was prepared by cloning the intron from pCINeo (Promega). The synthetic intron from pCI-Neo was amplified using standard PCR conditions and the primers:

CBProFpr174: 5'- AGCTGGCTAGCCACTGGGCAGGTAAGTATCA -3' and

CBProFpr175: 5'- TGGCGGGATCCTTAAGAGCTGTAATTGAACT -3'

resulting in a 332 bp PCR fragment. The fragment was cut with *NheI* and *BamHI* before
5 cloning into pCDNA3.1/HygR (obtained from Invitrogen) resulting in PF#34.

The expression cassette for hFVII was cloned between the *BamHI* and *HindIII* sites of PF#34, resulting in plasmid PF#226.

Example 3

10 *Expression of polypeptide variants in CHO K1 cells*

The cell line CHO K1 (ATCC # CCL-61) is seeded at 50% confluence in T-25 flasks using MEM α , 10% FCS (Gibco/BRL Cat # 10091), P/S and 5 μ g/ml phyloquinone and allowed to grow until confluent. The confluent mono cell layer is transfected with 5 μ g of the relevant plasmid described above using the Lipofectamine 2000 transfection agent (Life
15 technologies) according to the manufacturer's instructions. Twenty four hours post transfection a sample is drawn and quantified using *e.g.* an ELISA recognizing the EGF1 domain of hFVII. At this time point relevant selection (*e.g.* Hygromycin B) may be applied to the cells with the purpose of generating a pool of stable transfectants. When using CHO K1 cells and the Hygromycin B resistance gene as selectable marker on the plasmid, this is usually achieved
20 within one week.

Example 4

Generation of CHO K1 cells stably expressing polypeptide variants.

A vial of CHO-K1 transfectant pool is thawed and the cells seeded in a 175 cm² tissue
25 flask containing 25 ml of MEM α , 10% FCS, phyloquinone (5 μ g/ml), 100 U/l penicillin, 100 μ g/l streptomycin and grown for 24 hours. The cells are harvested, diluted and plated in 96 well microtiter plates at a cell density of 1/2-1 cell/well. After a week of growth, colonies of 20-100 cells are present in the wells and those wells containing only one colony are labelled. After a further two weeks, the media in all wells containing only one colony is substituted with 200
30 μ l fresh medium. After 24 hours, a medium sample is withdrawn and analysed by *e.g.* ELISA. High producing clones are selected and used to produce FVII or variant on large scale.

Example 5*Purification of polypeptide variants and subsequent activation*

FVII and FVII variants are purified as follows: The procedure is performed at 4°C. The harvested culture media from large-scale production is ultrafiltered using a Millipore TFF system with 30 KDa cut-off Pellicon membranes. After concentration of the medium, citrate is added to 5 mM and the pH is adjusted to 8.6. If necessary, the conductivity is lowered to below 10 mS/cm. Subsequently, the sample is applied to a Q-sepharose FF column, equilibrated with 50 mM NaCl, 10 mM Tris pH 8.6. After washing the column with 100 mM NaCl, 10 mM Tris pH 8.6, followed by 150 mM NaCl, 10 mM Tris pH 8.6, FVII is eluted using 10 mM Tris, 25 mM NaCl, 35 mM CaCl₂, pH 8.6.

For the second chromatographic step, an affinity column is prepared by coupling of a monoclonal Calcium-dependent antiGla-domain antibody to CNBr-activated Sepharose FF. About 5.5 mg antibody is coupled per ml resin. The column is equilibrated with 10 mM Tris, 100 mM NaCl, 35 mM CaCl₂, pH 7.5. NaCl is added to the sample to a concentration of 100 mM NaCl and the pH is adjusted to 7.4 -7.6. After O/N application of the sample, the column is washed with 100 mM NaCl, 35 mM CaCl₂, 10 mM Tris pH 7.5, and the FVII protein is eluted with 100 mM NaCl, 50 mM citrate, 75 mM Tris pH 7.5.

For the third chromatographic, the conductivity of the sample is lowered to below 10 mS/cm, if necessary, and the pH is adjusted to 8.6. The sample is then applied to a Q-sepharose column (equilibrated with 50 mM NaCl, 10 mM Tris pH 8.6) at a density around 3-5 mg protein per ml gel to obtain efficient activation. After application, the column is washed with 50 mM NaCl, 10 mM Tris pH 8.6 for about 4 hours with a flow of 3-4 column volumes (cv) per hour. The FVII protein is eluted using a gradient of 0-100% of 500 mM NaCl, 10 mM Tris pH 8.6 over 40 cv. FVII containing fractions are pooled.

For the final chromatographic step, the conductivity is lowered to below 10 mS/cm. Subsequently, the sample is applied to a Q-sepharose column (equilibrated with 140 mM NaCl, 10 mM glycylglycine pH 8.6) at a concentration of 3-5 mg protein per ml gel. The column is then washed with 140 mM NaCl, 10 mM glycylglycine pH 8.6 and FVII is eluted with 140 mM NaCl, 15 mM CaCl₂, 10 mM glycylglycine pH 8.6. The eluate is diluted to 10 mM CaCl₂ and the pH is adjusted 6.8-7.2. Finally, Tween-80 is added to 0.01% and the pH is adjusted to 5.5 for storage at -80°C.

CLAIMS

1. A Factor VII (FVII) or Factor VIIa (FVIIa) polypeptide variant having an amino acid sequence comprising 3-15 amino acid modifications relative to human Factor VII (hFVII) or
5 human Factor VIIa (hFVIIa) having the amino acid sequence shown in SEQ ID NO:2, wherein said amino acid sequence of the variant comprises an amino acid substitution in position 10 and 32 and wherein a sugar moiety is covalently attached to an introduced *in vivo* N-glycosylation site located outside the Gla domain.
- 10 2. The variant according to claim 1, wherein said substitution in position 10 is P10Q.
3. The variant according to claim 1 or 2, wherein said substitution in position 32 is K32E.
4. The variant according to any of the preceding claims, wherein said substitution in position 10
15 is P10Q and said substitution in position 32 is K32E.
5. The variant according to any of the preceding claims, wherein said variant comprises at least one further amino acid modification in the Gla domain.
- 20 6. The variant according to claim 5, wherein said further modification in the Gla domain comprises an amino acid substitution in position 33.
7. The variant according to claim 6, wherein a hydrophobic amino acid residue is introduced by substitution in position 33.
- 25 8. The variant according to claim 7, wherein said substitution is D33F.
9. The variant according to claim 8, wherein said variant comprises the substitutions P10Q+K32E+D33F.
- 30 10. The variant according to any of claims 5-9, wherein said further modification in the Gla domain comprises insertion of at least one amino acid residue between position 3 and 4.

11. The variant according to claim 10, wherein said further modification in the Gla domain comprises insertion of one amino acid residue between position 3 and 4.
12. The variant according to claim 10 or 11, wherein a hydrophobic amino acid residue is
5 inserted between position 3 and 4.
13. The variant according to claim 12, wherein said insertion is A3AY.
14. The variant according to claim 13, wherein said variant comprises the modifications
10 A3AY+P10Q+K32E.
15. The variant according to claim 13, wherein said variant comprises the modifications
A3AY+P10Q+K32E+D33F.
- 15 16. The variant according to any of claims 5-15, wherein said further modification in the Gla domain comprises an amino acid substitution in position 34.
17. The variant according to claim 16, wherein a negatively charged amino acid residue is introduced by substitution in position 34.
20
18. The variant according to claim 17, wherein said substitution is A34E.
19. The variant according to claim 18, wherein said variant comprises the substitutions
P10Q+K32E+A34E.
25
20. The variant according to claim 18, wherein said variant comprises the substitutions
P10Q+K32E+D33F+A34E.
21. The variant according to claim 18, wherein said variant comprises the modifications
30 A3AY+P10Q+K32E+A34E.
22. The variant according to claim 18, wherein said variant comprises the modifications
A3AY+P10Q+K32E+D33F+A34E.

23. The variant according to any of claims 5-22, wherein said further modification in the Gla domain comprises a substitution in a position selected from the group consisting of position 8, 11 and 28.

5 24. The variant according to claim 23, wherein said substitution in position 28 is R28F or R28E.

25. The variant according to any of the preceding claims, wherein no modifications are made in residues 6, 7, 14, 16, 19, 20, 25, 26, 29 and 35.

10

26. The variant according to any of the preceding claims, wherein said *in vivo* N-glycosylation site is introduced by substitution.

27. The variant according to claim 26, wherein said *in vivo* N-glycosylation site is introduced
15 into a position comprising an amino acid residue having at least 25% of its side chain exposed to the surface (as defined in Example 1 herein).

28. The variant according to claim 27, wherein said *in vivo* N-glycosylation site is introduced
20 into a position comprising an amino acid residue having at least 50% of its side chain exposed to the surface (as defined in Example 1 herein).

29. The variant according to any of claims 26-28, wherein said *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of A51N, G58N, T106N, K109N, G124N, K143N+N145T, A175T, I205S, I205T, V253N, T267N, T267N+S269T,
25 S314N+K316S, S314N+K316T, R315N+V317S, R315N+V317T, K316N+G318S, K316N+G318T, G318N, D334N and combinations thereof.

30. The variant according to claim 29, wherein said *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of A51N, G58N, T106N, K109N, G124N,
30 K143N+N145T, A175T, I205T, V253N, T267N+S269T, S314N+K316T, R315N+V317T, K316N+G318T, G318N, D334N and combinations thereof.

31. The variant according to claim 30, wherein said *in vivo* N-glycosylation site is introduced by a substitution selected from the group consisting of T106N, A175T, I205T, V253N, T267N+S269T and combinations thereof.

5 32. The variant according to any of claims 16-31, wherein one *in vivo* N-glycosylation site has been introduced by substitution.

33. The variant according to any of claims 16-31, wherein two or more *in vivo* N-glycosylation sites have been introduced by substitution.

10

34. The variant according to claim 33, wherein two *in vivo* N-glycosylation sites have been introduced by substitution.

35. The variant according to claim 33 or 34, wherein said *in vivo* N-glycosylation sites have
 15 been introduced by substitutions selected from the group consisting of A51N+G58N,
 A51N+T106N, A51N+K109N, A51N+G124N, A51N+K143N+N145T, A51N+A175T,
 A51N+I205T, A51N+V253N, A51N+T267N+S269T, A51N+S314N+K316T,
 A51N+R315N+V317T, A51N+K316N+G318T, A51N+G318N, A51N+D334N,
 G58N+T106N, G58N+K109N, G58N+G124N, G58N+K143N+N145T, G58N+A175T,
 20 G58N+I205T, G58N+V253N, G58N+T267N+S269T, G58N+S314N+K316T,
 G58N+R315N+V317T, G58N+K316N+G318T, G58N+G318N, G58N+D334N,
 T106N+K109N, T106N+G124N, T106N+K143N+N145T, T106N+A175T, T106N+I205T,
 T106N+V253N, T106N+T267N+S269T, T106N+S314N+K316T, T106N+R315N+V317T,
 T106N+K316N+G318T, T106N+G318N, T106N+D334N, K109N+G124N,
 25 K109N+K143N+N145T, K109N+A175T, K109N+I205T, K109N+V253N,
 K109N+T267N+S269T, K109N+S314N+K316T, K109N+R315N+V317T,
 K109N+K316N+G318T, K109N+G318N, K109N+D334N, G124N+K143N+N145T,
 G124N+A175T, G124N+I205T, G124N+V253N, G124N+T267N+S269T,
 G124N+S314N+K316T, G124N+R315N+V317T, G124N+K316N+G318T, G124N+G318N,
 30 G124N+D334N, K143N+N145T+A175T, K143N+N145T+I205T, K143N+N145T+V253N,
 K143N+N145T+T267N+S269T, K143N+N145T+S314N+K316T,
 K143N+N145T+R315N+V317T, K143N+N145T+K316N+G318T, K143N+N145T+G318N,
 K143N+N145T+D334N, A175T+I205T, A175T+V253N, A175T+T267N+S269T,
 A175T+S314N+K316T, A175T+R315N+V317T, A175T+K316N+G318T, A175T+G318N,

- A175T+D334N, I205T+V253N, I205T+T267N+S269T, I205T+S314N+K316T, I205T+R315N+V317T, I205T+K316N+G318T, I205T+G318N, I205T+D334N, V253N+T267N+S269T, V253N+S314N+K316T, V253N+R315N+V317T, V253N+K316N+G318T, V253N+G318N, V253N+D334N, T267N+S269T+S314N+K316T, T267N+S269T+R315N+V317T, T267N+S269T+K316N+G318T, T267N+S269T+G318N, T267N+S269T+D334N, S314N+K316T+R315N+V317T, S314N+K316T+G318N, S314N+K316T+D334N, R315N+V317T+K316N+G318T, R315N+V317T+G318N, R315N+V317T+D334N and G318N+D334N.
- 10 36. The variant according to claim 35, wherein said *in vivo* N-glycosylation sites have been introduced by substitutions selected from the group consisting of T106N+A175T, T106N+I205T, T106N+V253N, T106N+T267N+S269T, A175T+I205T, A175T+V253N, A175T+T267N+S269T, I205T+V253N, I205T+T267N+S269T and V253N+T267N+S269T.
- 15 37. The variant according to claim 36, wherein said *in vivo* N-glycosylation sites have been introduced by substitutions selected from the group consisting of T106N+I205T, T106N+V253N and I205T+T267N+S269T.
38. The variant according to any of claims 26-37, wherein three or more *in vivo* N-glycosylation sites have been introduced by substitution.
- 20 39. The variant according to claim 38, wherein three *in vivo* N-glycosylation sites have been introduced by substitution.
40. The variant according to claim 38 or 39, wherein said *in vivo* N-glycosylation sites have been introduced by substitutions selected from the group consisting of I205T+V253N+T267N+S269T and T106N+I205T+V253N.
41. The variant according to any of the preceding claims, wherein said variant is in its activated form.
- 30 42. The variant according to any of the preceding claims, wherein said variant, in its activated form, has at least 10% of the amidolytic activity of rhFVIIa when assayed in the "Amidolytic Assay" described herein.

43. The variant according to any of the preceding claims, wherein said variant, in its activated form, has at least 10% of the clotting activity of rhFVIIa when assayed in the "Clotting Assay" described herein.

5

44. A nucleotide sequence encoding a variant as defined in any of claims 1-43.

45. An expression vector comprising a nucleotide sequence as defined in claim 44.

10 46. A host cell comprising a nucleotide sequence as defined in claim 44 or an expression vector as defined in claim 45.

47. The host cell according to claim 46, wherein said host cell is a gammacarboxylating cell capable of *in vivo* glycosylation.

15

48. A pharmaceutical composition comprising a variant as defined in any of claims 1-43, and a pharmaceutical acceptable carrier or excipient.

49. A variant as defined in any of claims 1-43, or a pharmaceutical composition as defined in
20 claim 48, for use as a medicament.

50. Use of a variant as defined in any of claims 1-43 for the manufacture of a medicament for the treatment of a disease or a disorder wherein clot formation is desirable.

25 51. Use according to claim 50, wherein said disease or disorder is selected from the group consisting of hemorrhages, including brain hemorrhages, severe uncontrolled bleedings, such as trauma, bleedings in patients undergoing living transplantations, bleeding in patients undergoing resection and variceal bleedings.

30 52. Use according to claim 51, wherein said disease or disorder is trauma.

53. Use according to claim 52, wherein said disease or disorder is blunt trauma.

54. Use according to claim 52, wherein said disease or disorder is penetrative trauma.

55. A method for treating a mammal having a disease or a disorder wherein clot formation is desirable, comprising administering to a mammal in need thereof an effective amount of the variant as defined in any of claims 1-43 or the pharmaceutical composition as defined in claim
5 48.

56. The method according to claim 55, wherein said disease or disorder is selected from the group consisting of hemorrhages, including brain hemorrhages, severe uncontrolled bleedings, such as trauma, bleedings in patients undergoing living transplantations, bleeding in patients
10 undergoing resection and variceal bleedings.

57. The method according to claim 56, wherein said disease or disorder is trauma.

58. The method according to claim 57, wherein said disease or disorder is blunt trauma.
15

59. The method according to claim 57, wherein said disease or disorder is penetrative trauma.

SEQUENCE LISTING

<110> Maxygen ApS; Maxygen Holdings

<120> Factor VII or VIIa Polypeptide Variants

<130> 0259wo210

<140>

<141>

<160> 4

<170> PatentIn Ver. 2.1

<210> 1

<211> 1338

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (115)..(1335)

<400> 1

```

atgggtcagcc aggcctctcg cctcctgtgc ctgctcctgg ggctgcaggg ctgcctggct 60
gccgtcttcg tcaccagga ggaagcccat ggcgtcctgc atcgccggcg ccgg gcc 117
                                     Ala
                                     1

aat gcc ttt ctg gaa gag ctc cgc cct ggc tcc ctg gaa cgc gaa tgc 165
Asn Ala Phe Leu Glu Glu Leu Arg Pro Gly Ser Leu Glu Arg Glu Cys
      5              10              15

aaa gag gaa cag tgc agc ttt gag gaa gcc cgg gag att ttc aaa gac 213
Lys Glu Glu Gln Cys Ser Phe Glu Glu Ala Arg Glu Ile Phe Lys Asp
      20              25              30

gct gag cgg acc aaa ctg ttt tgg att agc tat agc gat ggc gat cag 261
Ala Glu Arg Thr Lys Leu Phe Trp Ile Ser Tyr Ser Asp Gly Asp Gln
      35              40              45

tgc gcc tcc agc cct tgc cag aac ggg ggc tcc tgc aaa gac cag ctg 309
Cys Ala Ser Ser Pro Cys Gln Asn Gly Gly Ser Cys Lys Asp Gln Leu
      50              55              60              65

cag agc tat atc tgc ttc tgc ctg cct gcc ttt gag ggg cgc aat tgc 357
Gln Ser Tyr Ile Cys Phe Cys Leu Pro Ala Phe Glu Gly Arg Asn Cys
      70              75              80

gaa acc cat aag gat gac cag ctg att tgc gtc aac gaa aac ggg ggc 405
Glu Thr His Lys Asp Asp Gln Leu Ile Cys Val Asn Glu Asn Gly Gly
      85              90              95

tgc gag cag tac tgc agc gat cac acg ggc acg aag cgg agc tgc cgc 453
Cys Glu Gln Tyr Cys Ser Asp His Thr Gly Thr Lys Arg Ser Cys Arg
      100              105              110

```

tgc cac gaa ggc tat agc ctc ctg gct gac ggg gtg tcc tgc acg ccc	501
Cys His Glu Gly Tyr Ser Leu Leu Ala Asp Gly Val Ser Cys Thr Pro	
115 120 125	
acg gtg gaa tac cct tgc ggg aag att ccc att cta gaa aag cgg aac	549
Thr Val Glu Tyr Pro Cys Gly Lys Ile Pro Ile Leu Glu Lys Arg Asn	
130 135 140 145	
gct agc aaa ccc cag ggc cgg atc gtc ggc ggg aag gtc tgc cct aag	597
Ala Ser Lys Pro Gln Gly Arg Ile Val Gly Gly Lys Val Cys Pro Lys	
150 155 160	
ggg gag tgc ccc tgg cag gtc ctg ctc ctg gtc aac ggg gcc cag ctg	645
Gly Glu Cys Pro Trp Gln Val Leu Leu Leu Val Asn Gly Ala Gln Leu	
165 170 175	
tgc ggc ggg acc ctc atc aat acc att tgg gtc gtg tcc gcc gct cac	693
Cys Gly Gly Thr Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala His	
180 185 190	
tgc ttc gat aag att aag aat tgg cgg aac ctc atc gct gtg ctc ggc	741
Cys Phe Asp Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu Gly	
195 200 205	
gaa cac gat ctg tcc gag cat gac ggg gac gaa cag tcc cgc cgg gtg	789
Glu His Asp Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg Val	
210 215 220 225	
gct cag gtc atc att ccc tcc acc tat gtg cct ggc acg acc aat cac	837
Ala Gln Val Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn His	
230 235 240	
gat atc gct ctg ctc cgc ctc cac cag ccc gtc gtg ctc acc gat cac	885
Asp Ile Ala Leu Leu Arg Leu His Gln Pro Val Val Leu Thr Asp His	
245 250 255	
gtc gtg cct ctg tgc ctg cct gag cgg acc ttt agc gaa cgc acg ctg	933
Val Val Pro Leu Cys Leu Pro Glu Arg Thr Phe Ser Glu Arg Thr Leu	
260 265 270	
gct ttc gtc cgc ttt agc ctc gtg tcc ggc tgg ggc cag ctg ctc gac	981
Ala Phe Val Arg Phe Ser Leu Val Ser Gly Trp Gly Gln Leu Leu Asp	
275 280 285	
cgg ggc gct acc gct ctc gag ctg atg gtg ctc aac gtc ccc cgg ctg	1029
Arg Gly Ala Thr Ala Leu Glu Leu Met Val Leu Asn Val Pro Arg Leu	
290 295 300 305	
atg acc cag gac tgc ctg cag cag tcc cgc aaa gtg ggg gac tcc ccc	1077
Met Thr Gln Asp Cys Leu Gln Gln Ser Arg Lys Val Gly Asp Ser Pro	
310 315 320	
aat atc acg gag tat atg ttt tgc gct ggc tat agc gat ggc tcc aag	1125
Asn Ile Thr Glu Tyr Met Phe Cys Ala Gly Tyr Ser Asp Gly Ser Lys	
325 330 335	

gat agc tgc aag ggg gac tcc ggc ggg ccc cat gcc acg cac tat cgc 1173
Asp Ser Cys Lys Gly Asp Ser Gly Gly Pro His Ala Thr His Tyr Arg
340 345 350

ggg acc tgg tac ctc acc ggg atc gtc agc tgg ggc cag ggc tgc gcc 1221
Gly Thr Trp Tyr Leu Thr Gly Ile Val Ser Trp Gly Gln Gly Cys Ala
355 360 365

acg gtg ggg cac ttt ggc gtc tac acg cgc gtc agc cag tac att gag 1269
Thr Val Gly His Phe Gly Val Tyr Thr Arg Val Ser Gln Tyr Ile Glu
370 375 380 385

tgg ctg cag aag ctc atg cgg agc gaa ccc cgg ccc ggg gtg ctc ctg 1317
Trp Leu Gln Lys Leu Met Arg Ser Glu Pro Arg Pro Gly Val Leu Leu
390 395 400

cgg gcc cct ttc cct tga taa 1338
Arg Ala Pro Phe Pro
405

<210> 2
<211> 406
<212> PRT
<213> Homo sapiens

<400> 2
Ala Asn Ala Phe Leu Glu Glu Leu Arg Pro Gly Ser Leu Glu Arg Glu
1 5 10 15
Cys Lys Glu Glu Gln Cys Ser Phe Glu Glu Ala Arg Glu Ile Phe Lys
20 25 30
Asp Ala Glu Arg Thr Lys Leu Phe Trp Ile Ser Tyr Ser Asp Gly Asp
35 40 45
Gln Cys Ala Ser Ser Pro Cys Gln Asn Gly Gly Ser Cys Lys Asp Gln
50 55 60
Leu Gln Ser Tyr Ile Cys Phe Cys Leu Pro Ala Phe Glu Gly Arg Asn
65 70 75 80
Cys Glu Thr His Lys Asp Asp Gln Leu Ile Cys Val Asn Glu Asn Gly
85 90 95
Gly Cys Glu Gln Tyr Cys Ser Asp His Thr Gly Thr Lys Arg Ser Cys
100 105 110
Arg Cys His Glu Gly Tyr Ser Leu Ala Asp Gly Val Ser Cys Thr
115 120 125
Pro Thr Val Glu Tyr Pro Cys Gly Lys Ile Pro Ile Leu Glu Lys Arg
130 135 140
Asn Ala Ser Lys Pro Gln Gly Arg Ile Val Gly Gly Lys Val Cys Pro
145 150 155 160
Lys Gly Glu Cys Pro Trp Gln Val Leu Leu Val Asn Gly Ala Gln
165 170 175
Leu Cys Gly Gly Thr Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala
180 185 190
His Cys Phe Asp Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu
195 200 205
Gly Glu His Asp Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg
210 215 220
Val Ala Gln Val Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn
225 230 235 240
His Asp Ile Ala Leu Leu Arg Leu His Gln Pro Val Val Leu Thr Asp
245 250 255

His	Val	Val	Pro	Leu	Cys	Leu	Pro	Glu	Arg	Thr	Phe	Ser	Glu	Arg	Thr
			260					265					270		
Leu	Ala	Phe	Val	Arg	Phe	Ser	Leu	Val	Ser	Gly	Trp	Gly	Gln	Leu	Leu
		275					280					285			
Asp	Arg	Gly	Ala	Thr	Ala	Leu	Glu	Leu	Met	Val	Leu	Asn	Val	Pro	Arg
		290					295				300				
Leu	Met	Thr	Gln	Asp	Cys	Leu	Gln	Gln	Ser	Arg	Lys	Val	Gly	Asp	Ser
305					310					315				320	
Pro	Asn	Ile	Thr	Glu	Tyr	Met	Phe	Cys	Ala	Gly	Tyr	Ser	Asp	Gly	Ser
				325					330					335	
Lys	Asp	Ser	Cys	Lys	Gly	Asp	Ser	Gly	Gly	Pro	His	Ala	Thr	His	Tyr
			340					345					350		
Arg	Gly	Thr	Trp	Tyr	Leu	Thr	Gly	Ile	Val	Ser	Trp	Gly	Gln	Gly	Cys
		355					360					365			
Ala	Thr	Val	Gly	His	Phe	Gly	Val	Tyr	Thr	Arg	Val	Ser	Gln	Tyr	Ile
		370				375					380				
Glu	Trp	Leu	Gln	Lys	Leu	Met	Arg	Ser	Glu	Pro	Arg	Pro	Gly	Val	Leu
385					390					395				400	
Leu	Arg	Ala	Pro	Phe	Pro										
				405											

<210> 3

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer
CBProFpr174

<400> 3

agctggctag ccactgggca ggtaagtatc a

31

<210> 4

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer
CBProFpr175

<400> 4

tggcgggatc cttaagagct gtaattgaac t

31

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00276

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/64 C07K14/745 A61K38/36 A61P7/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data, BIOSIS, EMBASE, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	AMIT M. SHAH ET AL : "Manipulation of the membrane binding site of vitamin K-dependent proteins: Enhanced biological function of human factor VII" PROC. NATL. ACAD. SCI., vol. 95, April 1998 (1998-04), pages 4229-4234, XP002254307 the whole document	1-59
Y	GARY L. NELSESTUEN ET AL: "Elevated Function of Blood Clotting Factor VIIa Mutants That Have Enhanced Affinity for Membranes" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 276, no. 43, October 2001 (2001-10), pages 39825-39831, XP002254308 the whole document	1-59

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 September 2003

Date of mailing of the international search report

30 SEP 2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

CAROLINA PALMCRANTZ/EÖ

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00276

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 01 58935 A (MAXYGEN APS) 16 August 2001 (2001-08-16) the whole document	1-59
A	--- WO 02 22776 A (NOVO NORDISK AS ; PERSSON EGON (DK); OLSEN OLE HVILSTED (DK)) 21 March 2002 (2002-03-21) page 18, line 31 -page 19, line 4 -----	1-59

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00276

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: **55-59**
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 55-59 relate to methods of treatment of the human or animal body by therapy (Rule 39.1. (iv)). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/DK 03/00276

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0158935	A	16-08-2001	AU 3153501 A	20-08-2001
			CA 2397347 A1	16-08-2001
			CN 1400910 T	05-03-2003
			WO 0158935 A2	16-08-2001
			EP 1257295 A2	20-11-2002
			JP 2003521930 T	22-07-2003
			NO 20023804 A	25-09-2002
			US 2003096338 A1	22-05-2003

WO 0222776	A	21-03-2002	AU 5614801 A	12-11-2001
			AU 8755001 A	26-03-2002
			CA 2420892 A1	21-03-2002
			CZ 20030611 A3	13-08-2003
			WO 0182943 A2	08-11-2001
			WO 0222776 A2	21-03-2002
			EP 1280548 A2	05-02-2003
			EP 1319067 A2	18-06-2003
			NO 20031129 A	12-05-2003
			US 2003100740 A1	29-05-2003
			US 2003104978 A1	05-06-2003
